

Examples from Multilevel Software Comparative Reviews

Douglas Bates
R Development Core Team
Douglas.Bates@R-project.org

January 21, 2005

Abstract

The Center for Multilevel Modelling at the Institute of Education, London maintains a web site of “Software reviews of multilevel modeling packages”. The data sets discussed in the reviews are available at this web site. We have incorporated these data sets in the `lme4` package for R and, in this vignette, provide the results of fitting several models to these data sets.

```
Loading required package: lme4
Loading required package: Matrix
Loading required package: latticeExtra
```

1 Introduction

2 Two-level normal models

The `Exam` data set is used in fitting examples of two-level normal multilevel models.

```
> str(Exam)
`data.frame':      4059 obs. of  10 variables:
 $ school  : Factor w/ 65 levels "1","2","3","4",...: 1 1 1 1 1 1 1 1 1 1 ...
 $ normexam: num   0.261  0.134 -1.724  0.968  0.544 ...
 $ schgend : Factor w/ 3 levels "mixed","boys",...: 1 1 1 1 1 1 1 1 1 1 ...
 $ schavg  : num   0.166  0.166  0.166  0.166  0.166 ...
 $ vr      : Factor w/ 3 levels "bottom 25%","mid 50%",...: 2 2 2 2 2 2 2 2 2 2 ...
 $ intake  : Factor w/ 3 levels "bottom 25%","mid 50%",...: 1 2 3 2 2 1 3 2 2 3 ...
```

```

$ standLRT: num 0.619 0.206 -1.365 0.206 0.371 ...
$ sex : Factor w/ 2 levels "F","M": 1 1 2 1 1 2 2 1 2 ...
$ type : Factor w/ 2 levels "Mxd","Sngl": 1 1 1 1 1 1 1 1 1 ...
$ student : Factor w/ 650 levels "1","2","3","4",...: 143 145 142 141 138 155 158 115 117 113 ...

> system.time(Eml <- lmer(normexam ~ standLRT + sex + schgend +
+ (1 | school), Exam), gc = TRUE)
[1] 0.13 0.00 0.13 0.00 0.00

> summary(Eml)

Linear mixed-effects model fit by REML
Formula: normexam ~ standLRT + sex + schgend + (1 | school)
Data: Exam
      AIC      BIC    logLik MLdeviance REMLdeviance
9361.673 9405.834 -4673.837  9325.501    9347.673
Random effects:
Groups Name Variance Std.Dev.
school (Intercept) 0.085829 0.29297
Residual 0.562534 0.75002
# of obs: 4059, groups: school, 65

Fixed effects:
              Estimate Std. Error DF t value Pr(>|t|)
(Intercept) -1.0493e-03 5.5569e-02 4054 -0.0189 0.98494
standLRT     5.5975e-01 1.2450e-02 4054 44.9601 < 2.2e-16
sexM         -1.6739e-01 3.4100e-02 4054 -4.9089 9.519e-07
schgendboys  1.7769e-01 1.1347e-01 4054 1.5659 0.11745
schgendgirls 1.5900e-01 8.9403e-02 4054 1.7784 0.07541

Correlation of Fixed Effects:
              (Intr) stnLRT sexM schgndb
standLRT     -0.014
sexM         -0.316 0.061
schgendboys  -0.395 -0.003 -0.145
schgendgrls  -0.622 0.009 0.197 0.245

```

There are some interesting aspects of data management that show up in the analysis of these data. The `student` variable is an identifier of the student within the `school`. It would be best to combine the indicators of school and student to get a unique identifier of the student.

```

> Exam$ids <- with(Exam, school:student)[, drop = TRUE]
> str(Exam)

`data.frame`:      4059 obs. of  11 variables:
 $ school : Factor w/ 65 levels "1","2","3","4",...: 1 1 1 1 1 1 1 1 1 ...
 $ normexam: num 0.261 0.134 -1.724 0.968 0.544 ...
 $ schgend : Factor w/ 3 levels "mixed","boys",...: 1 1 1 1 1 1 1 1 1 ...
 $ schavg  : num 0.166 0.166 0.166 0.166 0.166 ...
 $ vr      : Factor w/ 3 levels "bottom 25%","mid 50%",...: 2 2 2 2 2 2 2 2 2 ...
 $ intake  : Factor w/ 3 levels "bottom 25%","mid 50%",...: 1 2 3 2 2 1 3 2 2 3 ...
 $ standLRT: num 0.619 0.206 -1.365 0.206 0.371 ...
 $ sex     : Factor w/ 2 levels "F","M": 1 1 2 1 1 2 2 1 2 ...
 $ type    : Factor w/ 2 levels "Mxd","Sngl": 1 1 1 1 1 1 1 1 1 ...
 $ student : Factor w/ 650 levels "1","2","3","4",...: 143 145 142 141 138 155 158 115 117 113 ...
 $ ids     : Factor w/ 4055 levels "1:1","1:4","1:6",...: 48 49 47 46 45 50 51 39 40 38 ...

```

Notice that there are 4059 observations but only 4055 unique levels of student within school. We can check the ones that are duplicated

```
> Exam$ids[which(duplicated(Exam$ids))]  
[1] 43:86 50:39 52:2 52:21  
4055 Levels: 1:1 1:4 1:6 1:7 1:13 1:14 1:16 1:17 1:19 1:22 1:27 ... 65:155
```

One of these duplicated cases is particularly interesting. One of the students with the duplicated student id 86 in school 43 is the only male student in this mixed school. This is probably a case of a misrecorded school.

3 Three-level Normal Models

Data from the 1997 A-level Chemistry exam are available as Chem97.

```
> str(Chem97)  
`data.frame':      31022 obs. of  8 variables:  
 $ lea      : Factor w/ 131 levels "1","2","3","4",...: 1 1 1 1 1 1 1 1 1 ...  
 $ school   : Factor w/ 2410 levels "1","2","3","4",...: 1 1 1 1 1 1 1 1 1 ...  
 $ student  : Factor w/ 31022 levels "1","2","3","4",...: 1 2 3 4 5 6 7 8 9 10 ...  
 $ score    : num  4 10 10 10 8 10 6 8 4 10 ...  
 $ gender   : Factor w/ 2 levels "M","F": 2 2 2 2 2 2 2 2 2 2 ...  
 $ age      : num  3 -3 -4 -2 -1 4 1 4 3 0 ...  
 $ gcsecore: num  6.62 7.62 7.25 7.50 6.44 ...  
 $ gcsecnt  : num  0.339 1.339 0.964 1.214 0.158 ...  
  
> system.time(mC2 <- lmer(score ~ gcsecnt + (1 | school) +  
+ (1 | lea), Chem97), gc = TRUE)  
[1] 1.36 0.03 1.40 0.00 0.00  
  
> summary(mC2)  
  
Linear mixed-effects model fit by REML  
Formula: score ~ gcsecnt + (1 | school) + (1 | lea)  
Data: Chem97  
      AIC      BIC    logLik MLdeviance REMLdeviance  
141708.7 141750.4 -70849.33  141687.0    141698.7  
Random effects:  
Groups   Name      Variance  Std.Dev.  
school   (Intercept) 1.1709e+00 1.0821e+00  
lea      (Intercept) 5.1564e-10 2.2708e-05  
Residual                    5.1564e+00 2.2708e+00  
# of obs: 31022, groups: school, 2410; lea, 131  
  
Fixed effects:  
              Estimate Std. Error    DF t value Pr(>|t|)  
(Intercept) 5.6280e+00 2.7995e-02 31020  201.04 < 2.2e-16  
gcsecnt     2.4724e+00 1.6893e-02 31020   146.36 < 2.2e-16  
  
Correlation of Fixed Effects:  
      (Intr)  
gcsecnt 0.061
```

4 Two-level models for binary data

The data frame `Contraception` provides data from the Bangladesh fertility survey.

```
> str(Contraception)
`data.frame':      1934 obs. of  6 variables:
 $ woman   : Factor w/ 1934 levels "1","2","3","4",...: 1 2 3 4 5 6 7 8 9 10 ...
 $ district: Factor w/ 60 levels "1","2","3","4",...: 1 1 1 1 1 1 1 1 1 1 ...
 $ use      : Factor w/ 2 levels "N","Y": 1 1 1 1 1 1 1 1 1 1 ...
 $ livch    : Factor w/ 4 levels "0","1","2","3+": 4 1 3 4 1 1 4 4 2 4 ...
 $ age      : num   18.44 -5.56  1.44  8.44 -13.56 ...
 $ urban    : Factor w/ 2 levels "N","Y": 2 2 2 2 2 2 2 2 2 2 ...
> summary(Contraception[, -1])
      district      use      livch      age      urban
14      : 118   N:1175    0 :530   Min.    :-13.560000   N:1372
1       : 117   Y: 759    1 :356   1st Qu.: -7.559900   Y: 562
46      :  86                2 :305   Median : -1.559900
25      :  67                3+:743   Mean    :  0.002198
6       :  65                3rd Qu.:  6.440000
30      :  61                Max.    : 19.440000
(Other):1420
> system.time(mB1 <- GLMM(use ~ urban + age + livch, binomial,
+   Contraception, ~1 | district))
[1] 1.06 0.01 1.10 0.00 0.00
> summary(mB1)
Generalized Linear Mixed Model

Family: binomial family with logit link
Fixed: use ~ urban + age + livch
Data: Contraception
      AIC      BIC    logLik
2429.664 2474.203 -1206.832

Random effects:
      Groups      Name      Variance    Std.Dev.
district (Intercept)      0.21518      0.46387
# of obs: 1934, groups: district, 60

Estimated scale (compare to 1) 0.9844111

Fixed effects:
      Estimate Std. Error z value Pr(>|z|)
(Intercept) -1.6606460  0.1452147 -11.4358 < 2.2e-16
urbanY       0.7193097  0.1183317  6.0788 1.211e-09
age         -0.0261558  0.0078152 -3.3468 0.0008176
livch1       1.0921026  0.1565011  6.9782 2.989e-12
livch2       1.3545533  0.1729641  7.8314 4.824e-15
livch3+      1.3241531  0.1773558  7.4661 8.262e-14

Correlation of Fixed Effects:
      (Intr) urbanY age      livch1 livch2
urbanY -0.300
age      0.446 -0.046
livch1  -0.589  0.059 -0.211
livch2  -0.631  0.094 -0.378  0.488
livch3+ -0.748  0.098 -0.674  0.539  0.619
```

```

> system.time(mB2 <- GLMM(use ~ urban + age + livch, binomial,
+   Contraception, ~1 | district, method = "Laplace"))
[1] 35.51 0.10 35.67 0.00 0.00
> summary(mB2)
Generalized Linear Mixed Model

Family: binomial family with logit link
Fixed: use ~ urban + age + livch
Data: Contraception
      AIC      BIC    logLik
2417.616 2428.750 -1206.808

Random effects:
      Groups      Name      Variance  Std.Dev.
district (Intercept)      0.21239    0.46086
# of obs: 1934, groups: district, 60

Estimated scale (compare to 1) 0.9859619

Fixed effects:
      Estimate Std. Error z value Pr(>|z|)
(Intercept) -1.6897122  0.1459423 -11.5779 < 2.2e-16
urbanY       0.7329916  0.1192199  6.1482 7.835e-10
age          -0.0265952  0.0078781 -3.3758 0.000736
livchl       1.1091850  0.1576953  7.0337 2.011e-12
livch2       1.3763970  0.1743426  7.8948 2.908e-15
livch3+      1.3452373  0.1787274  7.5268 5.202e-14

Correlation of Fixed Effects:
      (Intr) urbanY age  livchl livch2
urbanY -0.301
age      0.448 -0.046
livchl  -0.589 0.059 -0.211
livch2  -0.631 0.094 -0.378 0.487
livch3+ -0.749 0.099 -0.674 0.538 0.618

> system.time(mB3 <- GLMM(use ~ urban + age + livch, family = binomial,
+   data = Contraception, random = ~urban | district))
[1] 1.44 0.01 1.44 0.00 0.00
> summary(mB3)
Generalized Linear Mixed Model

Family: binomial family with logit link
Fixed: use ~ urban + age + livch
Data: Contraception
      AIC      BIC    logLik
2225.720 2281.394 -1102.860

Random effects:
      Groups      Name      Variance Std.Dev. Corr
district (Intercept) 0.38774  0.62269
      urbanY      0.66745  0.81698 -0.793
# of obs: 1934, groups: district, 60

Estimated scale (compare to 1) 0.9759564

```

```
Fixed effects:
              Estimate Std. Error z value Pr(>|z|)
(Intercept) -1.6665200  0.1572532 -10.5977 < 2.2e-16
urbanY       0.7914232  0.1681257   4.7073 2.510e-06
age         -0.0258502  0.0079082  -3.2688 0.00108
livch1       1.0987723  0.1580051   6.9540 3.550e-12
livch2       1.3342511  0.1745854   7.6424 2.132e-14
livch3+      1.3227367  0.1795440   7.3672 1.743e-13
```

```
Correlation of Fixed Effects:
              (Intr) urbanY age   livch1 livch2
urbanY      -0.481
age          0.416 -0.036
livch1      -0.548  0.038 -0.211
livch2      -0.586  0.068 -0.378  0.487
livch3+     -0.695  0.062 -0.674  0.537  0.616
```

5 Growth curve model for repeated measures data

```
> str(Oxboys)
`data.frame`:      234 obs. of  4 variables:
 $ Subject : Factor w/ 26 levels "1","10","11",...: 1 1 1 1 1 1 1 1 1 12 ...
 $ age      : num  -1.0000 -0.7479 -0.4630 -0.1643 -0.0027 ...
 $ height   : num   140 143 145 147 148 ...
 $ Occasion: Factor w/ 9 levels "1","2","3","4",...: 1 2 3 4 5 6 7 8 9 1 ...
- attr(*, "ginfo")=List of 7
 ..$ formula      :Class 'formula' length 3 height ~ age | Subject
 .. ..- attr(*, ".Environment")=length 6 <environment>
 ..$ order.groups: logi TRUE
 ..$ FUN           :function (x)
 .. ..- attr(*, "source")= chr "function (x) max(x, na.rm = TRUE)"
 ..$ outer         : NULL
 ..$ inner         : NULL
 ..$ labels        :List of 2
 .. ..$ age        : chr "Centered age"
 .. ..$ height      : chr "Height"
 ..$ units         :List of 1
 .. ..$ height      : chr "(cm)"

> system.time(mX1 <- lmer(height ~ age + I(age^2) + I(age^3) +
+ I(age^4) + (age + I(age^2) | Subject), Oxboys), gc = TRUE)
[1] 0.39 0.00 0.39 0.00 0.00

> summary(mX1)
Linear mixed-effects model fit by REML
Formula: height ~ age + I(age^2) + I(age^3) + I(age^4) + (age + I(age^2) | Subject)
Data: Oxboys
      AIC      BIC    logLik MLdeviance REMLdeviance
651.9081 693.372 -313.9541  625.3593    627.9081

Random effects:
Groups   Name             Variance Std.Dev. Corr
Subject (Intercept) 64.03165 8.00198
          age        2.86409 1.69236  0.614
          I(age^2)   0.67428 0.82115  0.215 0.658
```

```

Residual                0.21738 0.46624
# of obs: 234, groups: Subject, 26

Fixed effects:
              Estimate Std. Error  DF t value  Pr(>|t|)
(Intercept) 149.01887    1.57033 229 94.8968 < 2.2e-16
age          6.17418     0.35650 229 17.3189 < 2.2e-16
I(age^2)     1.12823     0.35144 229  3.2103 0.001516
I(age^3)     0.45385     0.16246 229  2.7937 0.005653
I(age^4)    -0.37690     0.30018 229 -1.2556 0.210554

Correlation of Fixed Effects:
      (Intr) age    I(g^2) I(g^3)
age      0.572
I(age^2) 0.076 0.264
I(age^3) -0.001 -0.340 0.025
I(age^4) 0.021 0.016 -0.857 -0.021

> system.time(mX2 <- lmer(height ~ poly(age, 4) + (age + I(age^2) |
+   Subject), Oxboys), gc = TRUE)
[1] 0.38 0.00 0.38 0.00 0.00

> summary(mX2)

Linear mixed-effects model fit by REML
Formula: height ~ poly(age, 4) + (age + I(age^2) | Subject)
Data: Oxboys
      AIC      BIC    logLik MLdeviance REMLdeviance
640.8686 682.3324 -308.4343  625.3593    616.8686

Random effects:
Groups Name      Variance Std.Dev. Corr
Subject (Intercept) 64.03118 8.00195
      age          2.86407 1.69236 0.614
      I(age^2)     0.67428 0.82115 0.215 0.658
Residual                0.21738 0.46624
# of obs: 234, groups: Subject, 26

Fixed effects:
              Estimate Std. Error  DF t value  Pr(>|t|)
(Intercept) 149.51976    1.59026 229 94.0221 < 2.2e-16
poly(age, 4)1  64.54095    3.32781 229 19.3944 < 2.2e-16
poly(age, 4)2  4.20322    1.02361 229  4.1063 5.597e-05
poly(age, 4)3  1.29077    0.46628 229  2.7682 0.006098
poly(age, 4)4 -0.58547    0.46630 229 -1.2556 0.210554

Correlation of Fixed Effects:
      (Intr) p(,4)1 p(,4)2 p(,4)3
poly(ag,4)1 0.631
poly(ag,4)2 0.230 0.583
poly(ag,4)3 0.000 0.000 0.000
poly(ag,4)4 0.000 0.000 0.000 0.000

```

6 Cross-classification model

```
> str(ScotsSec)
```

```

`data.frame':      3435 obs. of  6 variables:
 $ verbal : num  11 0 -14 -6 -30 -17 -17 -11 -9 -19 ...
 $ attain : num  10 3 2 3 2 2 4 6 4 2 ...
 $ primary: Factor w/ 148 levels "1","2","3","4",...: 1 1 1 1 1 1 1 1 1 1 ...
 $ sex    : Factor w/ 2 levels "M","F": 1 2 1 1 2 2 2 1 1 1 ...
 $ social : num   0 0 0 20 0 0 0 0 0 0 ...
 $ second : Factor w/ 19 levels "1","2","3","4",...: 9 9 9 9 9 9 1 1 9 9 ...

> system.time(mS1 <- lmer(attain ~ sex + (1 | primary) + (1 |
+   second), ScotsSec), gc = TRUE)

[1] 0.32 0.00 0.31 0.00 0.00

> summary(mS1)

Linear mixed-effects model fit by REML
Formula: attain ~ sex + (1 | primary) + (1 | second)
Data: ScotsSec
      AIC      BIC    logLik MLdeviance REMLdeviance
17137.91 17168.62 -8563.956   17123.49     17127.91
Random effects:
Groups   Name      Variance Std.Dev.
primary (Intercept) 1.10962   1.0534
second  (Intercept) 0.36966   0.6080
Residual                    8.05511  2.8382
# of obs: 3435, groups: primary, 148; second, 19

Fixed effects:
              Estimate Std. Error   DF t value Pr(>|t|)
(Intercept) 5.2552e+00 1.8432e-01 3433  28.5107 < 2.2e-16
sexF        4.9851e-01 9.8255e-02 3433   5.0737 4.109e-07

Correlation of Fixed Effects:
      (Intr)
sexF -0.264

```