

Package ‘emplik’

October 13, 2022

Maintainer Mai Zhou <maizhou@gmail.com>

Version 1.2

Depends R (>= 3.0)

Imports quantreg, stats

Suggests KMsurv, boot

Author Mai Zhou. (Art Owen for el.test(). Yifan Yang for some C code.)

Description Empirical likelihood ratio tests for means/quantiles/hazards from possibly censored and/or truncated data. Now does regression too. This version contains some C code.

Title Empirical Likelihood Ratio for Censored/Truncated Data

License GPL (>= 2)

URL <http://www.ms.uky.edu/~mai/EmpLik.html>

NeedsCompilation yes

Repository CRAN

Date/Publication 2022-07-03 03:20:02 UTC

R topics documented:

emplik-package	2
BJoint	3
bjtest	4
bjtest1d	6
bjtestII	7
el.cen.EM	8
el.cen.EM2	11
el.cen.kmc1d	15
el.cen.test	18
el.ltrc.EM	20
el.test	22
el.test.wt	23
el.test.wt2	25

el.trun.test	27
emplikH.disc	28
emplikH.disc2	30
emplikH1.test	32
emplikH1B	34
emplikH1P	35
emplikH2.test	37
emplikH2B	39
emplikH2P	41
emplikHs.disc2	43
emplikHs.test2	45
findLnew	48
findUL	50
findUL2	52
findULold	54
findUnew	55
myeloma	57
RankRegTest	58
RankRegTestH	60
ROCnp	61
ROCnp2	63
smallcell	65
WRegEst	65
WRegTest	67

Index 69

emplik-package	<i>Empirical likelihood for mean functional/hazard functional with possibly censored data.</i>
----------------	--

Description

Empirical likelihood ratio tests and confidence intervals for means/hazards from possibly censored and/or truncated data. Now does regression too. The package contains some C code.

Details

```

Package: emplik
Type: Package
Version: 1.1-0
Date: 2020-05-18
License: GPL (>= 2)

```

For non-censored data and mean parameters, use `el.test()`.

For censored data and mean parameters, use `el.cen.EM2()`.

For censored data and hazard parameter, use `emplikH1.test()` [Poisson type likelihood]; use `emplikH.disc()` [binomial type likelihood].

For constructing confidence interval, use `findUL()`.

Author(s)

Mai Zhou (el.test is adapted from Owen's splus code. Yifan Yang for some C code.)

Maintainer: Mai Zhou <mai@ms.uky.edu> <maizhou@gmail.com>

References

Zhou, M. (2016). Empirical Likelihood Method in Survival Analysis. (Chapman and Hall/CRC Biostatistics Series) CRC press 2016

See Also

Another R package `kmc` for possible faster result for testing of mean with right censored data.

BJnoint

The Buckley-James censored regression estimator

Description

Compute the Buckley-James estimator in the regression model

$$y_i = \beta x_i + \epsilon_i$$

with right censored y_i . Iteration method.

Usage

```
BJnoint(x, y, delta, beta0 = NA, maxiter=30, error = 0.00001)
```

Arguments

<code>x</code>	a matrix or vector containing the covariate, one row per observation.
<code>y</code>	a numeric vector of length N, censored responses.
<code>delta</code>	a vector of length N, delta=0/1 for censored/uncensored.
<code>beta0</code>	an optional vector for starting value of iteration.
<code>maxiter</code>	an optional integer to control iterations.
<code>error</code>	an optional positive value to control iterations.

Details

This function compute the Buckley-James estimator when your model do not have an intercept term. Of course, if you include a column of 1's in the x matrix, it is also OK with this function and it is equivalent to having an intercept term. If your model do have an intercept term, then you could also (probably should) use the function `bj()` in the `Design` library. It should be more refined than `BJjoint` in the stopping rule for the iterations.

This function is included here mainly to produce the estimator value that may provide some useful information with the function `bjtest()`. For example you may want to test a beta value near the Buckley-James estimator.

Value

A list with the following components:

<code>beta</code>	the Buckley-James estimator.
<code>iteration</code>	number of iterations performed.

Author(s)

Mai Zhou.

References

Buckley, J. and James, I. (1979). Linear regression with censored data. *Biometrika*, **66** 429-36.

Examples

```
x <- matrix(c(rnorm(50,mean=1), rnorm(50,mean=2)), ncol=2,nrow=50)
## Suppose now we wish to test Ho: 2mu(1)-mu(2)=0, then
y <- 2*x[,1]-x[,2]
xx <- c(28,-44,29,30,26,27,22,23,33,16,24,29,24,40,21,31,34,-2,25,19)
```

`bjtest`

Test the Buckley-James estimator by Empirical Likelihood

Description

Use the empirical likelihood ratio and Wilks theorem to test if the regression coefficient is equal to beta.

The log empirical likelihood been maximized is

$$\sum_{d=1} \log \Delta F(e_i) + \sum_{d=0} \log[1 - F(e_i)];$$

where e_i are the residuals.

Usage

```
bjtest(y, d, x, beta)
```

Arguments

y a vector of length N, containing the censored responses.
d a vector (length N) of either 1's or 0's. d=1 means y is uncensored; d=0 means y is right censored.
x a matrix of size N by q.
beta a vector of length q. The value of the regression coefficient to be tested in the model $y_i = \beta x_i + \epsilon_i$

Details

The above likelihood should be understood as the likelihood of the error term, so in the regression model the error epsilon should be iid.

This version can handle the model where beta is a vector (of length q).

The estimation equations used when maximize the empirical likelihood is

$$0 = \sum d_i \Delta F(e_i)(x \cdot m[,i]) / (nw_i)$$

which was described in detail in the reference below.

Value

A list with the following components:

"-2LLR" the -2 loglikelihood ratio; have approximate chisq distribution under H_0 .
loge12 the log empirical likelihood, under estimating equation.
loge1 the log empirical likelihood of the Kaplan-Meier of e's.
prob the probabilities that max the empirical likelihood under estimating equation.

Author(s)

Mai Zhou.

References

Buckley, J. and James, I. (1979). Linear regression with censored data. *Biometrika*, **66** 429-36.
Zhou, M. and Li, G. (2008). Empirical likelihood analysis of the Buckley-James estimator. *Journal of Multivariate Analysis* **99**, 649-664.
Zhou, M. (2016) Empirical Likelihood Method in Survival Analysis. CRC Press.

Examples

```
xx <- c(28, -44, 29, 30, 26, 27, 22, 23, 33, 16, 24, 29, 24, 40, 21, 31, 34, -2, 25, 19)
```

bjtest1d

*Test the Buckley-James estimator by Empirical Likelihood, 1-dim only***Description**

Use the empirical likelihood ratio and Wilks theorem to test if the regression coefficient is equal to beta. For 1-dim beta only.

The log empirical likelihood been maximized is

$$\sum_{d=1} \log \Delta F(e_i) + \sum_{d=0} \log[1 - F(e_i)].$$

Usage

```
bjtest1d(y, d, x, beta)
```

Arguments

y	a vector of length N, containing the censored responses.
d	a vector of either 1's or 0's. d=1 means y is uncensored. d=0 means y is right censored.
x	a vector of length N, covariate.
beta	a number. the regression coefficient to be tested in the model $y = x \text{ beta} + \text{epsilon}$

Details

In the above likelihood, $e_i = y_i - x * \text{beta}$ is the residuals.

Similar to `bjtest()`, but only for 1-dim beta.

Value

A list with the following components:

"-2LLR"	the -2 loglikelihood ratio; have approximate chi square distribution under H_0 .
logel2	the log empirical likelihood, under estimating equation.
logel	the log empirical likelihood of the Kaplan-Meier of e's.
prob	the probabilities that max the empirical likelihood under estimating equation constraint.

Author(s)

Mai Zhou.

References

- Buckley, J. and James, I. (1979). Linear regression with censored data. *Biometrika*, **66** 429-36.
- Owen, A. (1990). Empirical likelihood ratio confidence regions. *Ann. Statist.* **18** 90-120.
- Zhou, M. and Li, G. (2008). Empirical likelihood analysis of the Buckley-James estimator. *Journal of Multivariate Analysis.* 649-664.

Examples

```
xx <- c(28, -44, 29, 30, 26, 27, 22, 23, 33, 16, 24, 29, 24, 40, 21, 31, 34, -2, 25, 19)
```

bjtestII	<i>Alternative test of the Buckley-James estimator by Empirical Likelihood</i>
----------	--

Description

Use the empirical likelihood ratio (alternative form) and Wilks theorem to test if the regression coefficient is equal to beta, based on the estimating equations.

The log empirical likelihood been maximized is

$$\sum_{j=1}^n \log p_j; \quad \sum p_j = 1$$

where the probability p_j is for the j th martingale differences of the estimating equations.

Usage

```
bjtestII(y, d, x, beta)
```

Arguments

- | | |
|------|--|
| y | a vector of length N, containing the censored responses. |
| d | a vector of length N. Either 1's or 0's. d=1 means y is uncensored; d=0 means y is right censored. |
| x | a matrix of size N by q. |
| beta | a vector of length q. The value of the regression coefficient to be tested in the model $Y_i = \beta x_i + \epsilon_i$ |

Details

The above likelihood should be understood as the likelihood of the martingale difference terms. For the definition of the Buckley-James martingale or estimating equation, please see the (2015) book in the reference list.

The estimation equations used when maximize the empirical likelihood is

$$0 = \sum d_i \Delta F(e_i)(x \cdot m[, i]) / (nw_i)$$

where e_i is the residuals, other details are described in the reference book of 2015 below.

The final test is carried out by `el.test`. So the output is similar to the output of `el.test`.

Value

A list with the following components:

"-2LLR"	the -2 loglikelihood ratio; have approximate chisq distribution under H_0 .
logel2	the log empirical likelihood, under estimating equation.
logel	the log empirical likelihood of the Kaplan-Meier of e's.
prob	the probabilities that max the empirical likelihood under estimating equation.

Author(s)

Mai Zhou.

References

- Zhou, M. (2016) Empirical Likelihood Methods in Survival Analysis. CRC Press.
- Buckley, J. and James, I. (1979). Linear regression with censored data. *Biometrika*, **66** 429-36.
- Zhou, M. and Li, G. (2008). Empirical likelihood analysis of the Buckley-James estimator. *Journal of Multivariate Analysis*, **99**, 649–664.
- Zhu, H. (2007) Smoothed Empirical Likelihood for Quantiles and Some Variations/Extension of Empirical Likelihood for Buckley-James Estimator, Ph.D. dissertation, University of Kentucky.

Examples

```
data(myeloma)
bjtestII(y=myeloma[,1], d=myeloma[,2], x=cbind(1, myeloma[,3]), beta=c(37, -3.4))
```

el.cen.EM	<i>Empirical likelihood ratio for mean with right, left or doubly censored data, by EM algorithm</i>
-----------	--

Description

This program uses EM algorithm to compute the maximized (wrt p_i) empirical log likelihood function for right, left or doubly censored data with the MEAN constraint:

$$\sum_{d_i=1} p_i f(x_i) = \int f(t) dF(t) = \mu.$$

Where $p_i = \Delta F(x_i)$ is a probability, d_i is the censoring indicator, 1(uncensored), 0(right censored), 2(left censored). It also returns those p_i .

The empirical log likelihood been maximized is

$$\sum_{d_i=1} \log \Delta F(x_i) + \sum_{d_i=0} \log[1 - F(x_i)] + \sum_{d_i=2} \log F(x_i).$$

Usage

```
e1.cen.EM(x,d,wt=rep(1,length(d)),fun=function(t){t},mu,maxit=50,error=1e-9,...)
```

Arguments

<code>x</code>	a vector containing the observed survival times.
<code>d</code>	a vector containing the censoring indicators, 1-uncensored; 0-right censored; 2-left censored.
<code>wt</code>	a weight vector (case weight). positive. same length as <code>d</code>
<code>fun</code>	a left continuous (weight) function used to calculate the mean as in H_0 . <code>fun(t)</code> must be able to take a vector input <code>t</code> . Default to the identity function $f(t) = t$.
<code>mu</code>	a real number used in the constraint, the mean value of $f(X)$.
<code>maxit</code>	an optional integer, used to control maximum number of iterations.
<code>error</code>	an optional positive real number specifying the tolerance of iteration error. This is the bound of the L_1 norm of the difference of two successive weights.
<code>...</code>	additional arguments, if any, to pass to <code>fun</code> .

Details

This implementation is all in R and have several for-loops in it. A faster version would use C to do the for-loop part. But this version seems faster enough and is easier to port to Splus.

We return the log likelihood all the time. Sometimes, (for right censored and no censor case) we also return the -2 log likelihood ratio. In other cases, you have to plot a curve with many values of the parameter, μ , to find out where is the place the log likelihood becomes maximum. And from there you can get -2 log likelihood ratio between the maximum location and your current parameter in H_0 .

In order to get a proper distribution as NPMLE, we automatically change the d for the largest observation to 1 (even if it is right censored), similar for the left censored, smallest observation. μ is a given constant. When the given constants μ is too far away from the NPMLE, there will be no distribution satisfy the constraint. In this case the computation will stop. The -2 Log empirical likelihood ratio should be infinite.

The constant μ must be inside $(\min f(x_i), \max f(x_i))$ for the computation to continue. It is always true that the NPMLE values are feasible. So when the computation stops, try move the μ closer to the NPMLE —

$$\sum_{d_i=1} p_i^0 f(x_i)$$

p_i^0 taken to be the jumps of the NPMLE of CDF. Or use a different `fun`.

Difference to the function `e1.cen.EM2`: here duplicate (input) observations are collapsed (with weight 2, 3, ... etc.) but those will stay separate by default in the `e1.cen.EM2`. This will lead to a different `loglik` value. But the -2LLR value should be same in either version.

Value

A list with the following components:

loglik	the maximized empirical log likelihood under the constraint. This may be different from the result of el.cen.EM2 because here the tied observations are collapsed into 1 with weight. (while el.cen.EM2 do not). However, the -2LLR should be the same.
times	locations of CDF that have positive mass. tied obs. are collapsed
prob	the jump size of CDF at those locations.
"-2LLR"	If available, it is Minus two times the Empirical Log Likelihood Ratio. Should be approximately chi-square distributed under H_0 .
Pval	The P-value of the test, using chi-square approximation.
lam	The Lagrange multiplier. Added 5/2007.

Author(s)

Mai Zhou

References

Zhou, M. (2005). Empirical likelihood ratio with arbitrary censored/truncated data by EM algorithm. *Journal of Computational and Graphical Statistics*, 643-656.

Murphy, S. and van der Vaart (1997) Semiparametric likelihood ratio inference. *Ann. Statist.* **25**, 1471-1509.

Examples

```
## example with tied observations
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
el.cen.EM(x,d,mu=3.5)
## we should get "-2LLR" = 1.2466...
myfun5 <- function(x, theta, eps) {
  u <- (x-theta)*sqrt(5)/eps
  INDE <- (u < sqrt(5)) & (u > -sqrt(5))
  u[u >= sqrt(5)] <- 0
  u[u <= -sqrt(5)] <- 1
  y <- 0.5 - (u - (u)^3/15)*3/(4*sqrt(5))
  u[ INDE ] <- y[ INDE ]
  return(u)
}
el.cen.EM(x, d, fun=myfun5, mu=0.5, theta=3.5, eps=0.1)
## example of using wt in the input. Since the x-vector contain
## two 5 (both d=1), and two 2(both d=0), we can also do
xx <- c(1, 1.5, 2, 3, 4, 5, 6, 4, 1, 4.5)
dd <- c(1, 1, 0, 1, 0, 1, 1, 1, 0, 1)
wt <- c(1, 1, 2, 1, 1, 2, 1, 1, 1, 1)
el.cen.EM(x=xx, d=dd, wt=wt, mu=3.5)
## this should be the same as the first example.
```

e1.cen.EM2

Empirical likelihood ratio test for a vector of means with right, left or doubly censored data, by EM algorithm

Description

This function is similar to `e1.cen.EM()`, but for multiple constraints. In the input there is a vector of observations $x = (x_1, \dots, x_n)$ and a function `fun`. The function `fun` should return the (n by k) matrix

$$(f_1(x), f_2(x), \dots, f_k(x)).$$

Also, the ordering of the observations, when consider censoring or redistributing-to-the-right, is according to the value of x , not `fun(x)`. So the probability distribution is for values x . This program uses EM algorithm to maximize (wrt p_i) empirical log likelihood function for right, left or doubly censored data with the MEAN constraint:

$$j = 1, 2, \dots, k \quad \sum_{d_i=1} p_i f_j(x_i) = \int f_j(t) dF(t) = \mu_j.$$

Where $p_i = \Delta F(x_i)$ is a probability, d_i is the censoring indicator, 1(uncensored), 0(right censored), 2(left censored). It also returns those p_i . The log likelihood function is defined as

$$\sum_{d_i=1} \log \Delta F(x_i) + \sum_{d_i=2} \log F(x_i) + \sum_{d_i=0} \log[1 - F(x_i)].$$

Usage

```
e1.cen.EM2(x,d,xc=1:length(x),fun,mu,maxit=50,error=1e-9,...)
```

Arguments

<code>x</code>	a vector containing the observed survival times.
<code>d</code>	a vector containing the censoring indicators, 1-uncensored; 0-right censored; 2-left censored.
<code>xc</code>	an optional vector of collapsing control values. If <code>xc[i]</code> <code>xc[j]</code> have different values then <code>(x[i], d[i])</code> , <code>(x[j], d[j])</code> will not merge into one observation with weight two, even if they are identical. Default is not to merge.
<code>fun</code>	a left continuous (weight) function that returns a matrix. The columns (=k) of the matrix is used to calculate the means and will be tested in H_0 . <code>fun(t)</code> must be able to take a vector input <code>t</code> .
<code>mu</code>	a vector of length k. Used in the constraint, as the mean of $f(X)$.
<code>maxit</code>	an optional integer, used to control maximum number of iterations.
<code>error</code>	an optional positive real number specifying the tolerance of iteration error. This is the bound of the L_1 norm of the difference of two successive weights.
<code>...</code>	additional inputs to pass to <code>fun()</code> .

Details

This implementation is all in R and have several for-loops in it. A faster version would use C to do the for-loop part. (but this version is easier to port to Splus, and seems faster enough).

We return the log likelihood all the time. Sometimes, (for right censored and no censor case) we also return the -2 log likelihood ratio. In other cases, you have to plot a curve with many values of the parameter, μ , to find out where the log likelihood becomes maximum. And from there you can get -2 log likelihood ratio between the maximum location and your current parameter in H_0 .

In order to get a proper distribution as NPMLE, we automatically change the d for the largest observation to 1 (even if it is right censored), similar for the left censored, smallest observation. μ is a given constant vector. When the given constants μ is too far away from the NPMLE, there will be no distribution satisfy the constraint. In this case the computation will stop. The -2 Log empirical likelihood ratio should be infinite.

The constant vector μ must be inside $(\min f(x_i), \max f(x_i))$ for the computation to continue. It is always true that the NPMLE values are feasible. So when the computation stops, try move the μ closer to the NPMLE —

$$\hat{\mu}_j = \sum_{d_i=1} p_i^0 f_j(x_i)$$

where p_i^0 taken to be the jumps of the NPMLE of CDF. Or use a different fun.

Difference to the function `el.cen.EM`: due to the introduction of input `xc` here in this function, the output `loglik` may be different compared to the function `el.cen.EM` due to not collapsing of duplicated input survival values. The -2LLR should be the same from both functions.

Value

A list with the following components:

<code>loglik</code>	the maximized empirical log likelihood under the constraints.
<code>times</code>	locations of CDF that have positive mass.
<code>prob</code>	the jump size of CDF at those locations.
<code>"-2LLR"</code>	If available, it is Minus two times the Empirical Log Likelihood Ratio. Should be approx. chi-square distributed under H_0 .
<code>Pval</code>	If available, the P-value of the test, using chi-square approximation.
<code>lam</code>	the Lagrange multiplier in the final EM step. (the M-step)

Author(s)

Mai Zhou

References

- Zhou, M. (2005). Empirical likelihood ratio with arbitrary censored/truncated data by EM algorithm. *Journal of Computational and Graphical Statistics*, 643-656.
- Zhou, M. (2002). Computing censored empirical likelihood ratio by EM algorithm. *Tech Report, Univ. of Kentucky, Dept of Statistics*

Examples

```

## censored regression with one right censored observation.
## we check the estimation equation, with the MLE inside myfun7.
y <- c(3, 5.3, 6.4, 9.1, 14.1, 15.4, 18.1, 15.3, 14, 5.8, 7.3, 14.4)
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 0)
### first we estimate beta, the MLE
lm.wfit(x=cbind(rep(1,12),x), y=y, w=WKM(x=y, d=d)$jump[rank(y)])$coef
## you should get 1.392885 and 2.845658
## then define myfun7 with the MLE value
myfun7 <- function(y, xmat) {
temp1 <- y - ( 1.392885 + 2.845658 * xmat)
return( cbind( temp1, xmat*temp1 )
}
## now test
el.cen.EM2(y,d, fun=myfun7, mu=c(0,0), xmat=x)
## we should get, Pval = 1 , as the MLE should.
## for other values of (a, b) inside myfun7, you get other Pval
##
rqfun1 <- function(y, xmat, beta, tau = 0.5) {
temp1 <- tau - (1-myfun55(y-beta*xmat))
return(xmat * temp1)
}
myfun55 <- function(x, eps=0.001){
u <- x*sqrt(5)/eps
INDE <- (u < sqrt(5)) & (u > -sqrt(5))
u[u >= sqrt(5)] <- 0
u[u <= -sqrt(5)] <- 1
y <- 0.5 - (u - (u)^3/15)*3/(4*sqrt(5))
u[ INDE ] <- y[ INDE ]
return(u)
}
## myfun55 is a smoothed indicator fn.
## eps should be between (1/sqrt(n), 1/n^0.75) [Chen and Hall]
el.cen.EM2(x=y,d=d,xc=1:12,fun=rqfun1,mu=0,xmat=x,beta=3.08,tau=0.44769875)
## default tau=0.5
el.cen.EM2(x=y,d=d,xc=1:12,fun=rqfun1,mu=0,xmat=x,beta=3.0799107404)
#####
### next 2 examples are testing the mean/median residual time
#####
mygfun <- function(s, age, muage) {as.numeric(s >= age)*(s-(age+muage))}
mygfun2 <- function(s, age, Mdage)
{as.numeric(s <= (age+Mdage)) - 0.5*as.numeric(s <= age)}
## Not run:
library(survival)
time <- cancer$time
status <- cancer$status-1
####for mean residual time
el.cen.EM2(x=time, d=status, fun=mygfun, mu=0, age=365.25, muage=234)$Pval
el.cen.EM2(x=time, d=status, fun=mygfun, mu=0, age=365.25, muage=323)$Pval
### for median residual time
el.cen.EM2(x=time, d=status, fun=mygfun2, mu=0.5, age=365.25, Mdage=184)$Pval

```

```

el.cen.EM2(x=time, d=status, fun=mygfun2, mu=0.5, age=365.25, Mdage=321)$Pval

## End(Not run)
## Not run:
#### For right censor only data (Kaplan-Meier) we can use this function to get a faster computation
#### by calling the kmc 0.2-2 package.
el.cen.R <- function (x, d, xc = 1:length(x), fun, mu, error = 1e-09, ...)
{
  xvec <- as.vector(x)
  d <- as.vector(d)
  mu <- as.vector(mu)
  xc <- as.vector(xc)
  n <- length(d)
  if (length(xvec) != n)
    stop("length of d and x must agree")
  if (length(xc) != n)
    stop("length of xc and d must agree")
  if (n <= 2 * length(mu) + 1)
    stop("Need more observations")
  if (any((d != 0) & (d != 1) ))
    stop("d must be 0(right-censored) or 1(uncensored)")
  if (!is.numeric(xvec))
    stop("x must be numeric")
  if (!is.numeric(mu))
    stop("mu must be numeric")

  funx <- as.matrix(fun(xvec, ...))
  pp <- ncol(funx)
  if (length(mu) != pp)
    stop("length of mu and ncol of fun(x) must agree")
  temp <- Wdataclean5(z = xvec, d, zc = xc, xmat = funx)
  x <- temp$value
  d <- temp$dd
  w <- temp$weight
  funx <- temp$xxmat
  d[length(d)] <- 1
  xd1 <- x[d == 1]
  if (length(xd1) <= 1)
    stop("need more distinct uncensored obs.")
  funxd1 <- funx[d == 1, ]
  xd0 <- x[d == 0]
  wd1 <- w[d == 1]
  wd0 <- w[d == 0]
  m <- length(xd0)

  pnw <- NA
  num <- NA
  if (m > 0) {
    gfun <- function(x) { return( fun(x, ...) - mu ) }
    temp <- kmc.solve(x=x, d=d, g=list(gfun))
    logel <- temp$loglik.h0
    logel00 <- temp$loglik.null
    lam <- - temp$lambd
  }
}

```

```

}
if (m == 0) {
num <- 0
temp6 <- el.test.wt2(x = funxd1, wt = wd1, mu)
pnew <- temp6$prob
lam <- temp6$lambda
logel <- sum(wd1 * log(pnew))
logel00 <- sum(wd1 * log(wd1/sum(wd1)))
}
tval <- 2 * (logel00 - logel)
list(loglik = logel, times = xd1, prob = pnew, lam = lam,
iters = num, `-2LLR` = tval, Pval = 1 - pchisq(tval,
df = length(mu)))
}

## End(Not run)

```

el.cen.kmc1d	<i>Empirical likelihood ratio for 1 mean constraint with right censored data</i>
--------------	--

Description

This program uses a fast recursive formula to compute the maximized (wrt p_i) empirical log likelihood ratio for right censored data with one MEAN constraint:

$$\sum_{d_i=1} p_i f(x_i) = \int f(t) dF(t) = \mu.$$

Where $p_i = \Delta F(x_i)$ is a probability, d_i is the censoring indicator, 1(uncensored), 0(right censored). It also returns those p_i .

The empirical log likelihood been maximized is

$$\sum_{d_i=1} \log \Delta F(x_i) + \sum_{d_i=0} \log[1 - F(x_i)].$$

Usage

```
el.cen.kmc1d(x, d, fun, mu, tol = .Machine$double.eps^0.5, step=0.001, ...)
```

Arguments

x	a vector containing the observed survival times.
d	a vector containing the censoring indicators, 1-uncensored; 0-right censored.
fun	a left continuous (weight) function used to calculate the mean as in H_0 . fun(t) must be able to take a vector input t.
mu	a real number used in the constraint, the mean value of $f(X)$.

tol	a small positive number, for the uniroot error tol.
step	a small positive number, for use in the uniroot function (as interval) to find lambda root. Sometimes uniroot will find the wrong root or no root, resulting a negative "-2LLR" or NA. Change the step to a different value often can fix this (but not always). Another sign of wrong root is that the sum of probabilities not sum to one, or has negative probability values.
...	additional arguments, if any, to pass to fun.

Details

This function is similar to the function in package `kmc`, but much simpler, i.e. all implemented in R and only for one mean. This implementation have two for-loops in R. A faster version would use C to do the for-loop part. But this version seems fast enough and is easier to port to Splus.

We return the log likelihood all the time. Sometimes, (for right censored case) we also return the -2 log likelihood ratio. In other cases, you have to plot a curve with many values of the parameter, μ , to find out where is the place the log likelihood becomes maximum. And from there you can get -2 log likelihood ratio between the maximum location and your current parameter in H_0 .

The input `step` is used in `uniroot` function to find a root of lambda. Sometimes a step value may lead to no root or result in a wrong root. You may try several values for the step to see. If the probabilities returned do not sum to one, then the lambda root is a wrong root. We want the root closest to zero.

In order to get a proper distribution as NPMLE, we automatically change the d for the largest observation to 1 (even if it is right censored). μ is a given constant. When the given constants μ is too far away from the NPMLE, there will be no distribution satisfy the constraint. In this case the computation will stop or return something ridiculous, (as negative -2LLR). The -2 Log empirical likelihood ratio may be +infinite.

The constant μ must be inside $(\min f(x_i), \max f(x_i))$ (with uncensored x_i) for the computation to continue. It is always true that the NPMLE values are feasible. So when the computation stops, try move the μ closer to the NPMLE —

$$\sum_{d_i=1} p_i^0 f(x_i)$$

p_i^0 taken to be the jumps of the NPMLE of CDF. Or use a different fun.

Value

A list with the following components:

loglik	the maximized empirical log likelihood under the constraint. Note, here the tied observations are not collapsed into one obs. with weight 2 (as in <code>el.cen.EM</code>), so the value may differ from those that do collapse the tied obs. In any case, the -2LLR should not differ (whether collaps or not).
times	locations of CDF that have positive mass.
prob	the jump size of CDF at those locations.

"-2LLR"	If available, it is minus two times the empirical Log Likelihood Ratio. Should be approximately chi-square distributed under Ho. If you got NA or negative value, then something is wrong, most likely the uniroot has found the wrong root. Suggest: use el.cen.EM2() which uses EM algorithm. It is more stable but slower.
Pval	The P-value of the test, using chi-square approximation.
lam	The Lagrange multiplier.

Author(s)

Mai Zhou

References

Zhou, M. and Yang, Y. (2015). A recursive formula for the Kaplan-Meier estimator with mean constraints and its application to empirical likelihood. *Computational Statistics* Vol. 30, Issue 4 pp. 1097-1109.

Zhou, M. (2005). Empirical likelihood ratio with arbitrary censored/truncated data by EM algorithm. *Journal of Computational and Graphical Statistics*, 14(3), 643-656.

Examples

```
x <- c(1, 1.5, 2, 3, 4.2, 5, 6.1, 5.3, 4.5, 0.9, 2.1, 4.3)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
ff <- function(x) {
  x - 3.7
}
el.cen.kmc1d(x=x, d=d, fun=ff, mu=0)
#####
## example with tied observations
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
el.cen.EM(x,d,mu=3.5)
## we should get "-2LLR" = 1.2466...
myfun5 <- function(x, theta, eps) {
  u <- (x-theta)*sqrt(5)/eps
  INDE <- (u < sqrt(5)) & (u > -sqrt(5))
  u[u >= sqrt(5)] <- 0
  u[u <= -sqrt(5)] <- 1
  y <- 0.5 - (u - (u)^3/15)*3/(4*sqrt(5))
  u[ INDE ] <- y[ INDE ]
  return(u)
}
el.cen.EM(x, d, fun=myfun5, mu=0.5, theta=3.5, eps=0.1)
## example of using wt in the input. Since the x-vector contain
## two 5 (both d=1), and two 2(both d=0), we can also do
xx <- c(1, 1.5, 2, 3, 4, 5, 6, 4, 1, 4.5)
dd <- c(1, 1, 0, 1, 0, 1, 1, 1, 0, 1)
wt <- c(1, 1, 2, 1, 1, 2, 1, 1, 1, 1)
el.cen.EM(x=xx, d=dd, wt=wt, mu=3.5)
## this should be the same as the first example.
```

 el.cen.test

 Empirical likelihood ratio for mean with right censored data, by QP.

Description

This program computes the maximized (wrt p_i) empirical log likelihood function for right censored data with the MEAN constraint:

$$\sum_i [d_i p_i g(x_i)] = \int g(t) dF(t) = \mu$$

where $p_i = \Delta F(x_i)$ is a probability, d_i is the censoring indicator. The d for the largest observation is always taken to be 1. It then computes the $-2 \log$ empirical likelihood ratio which should be approximately chi-square distributed if the constraint is true. Here $F(t)$ is the (unknown) CDF; $g(t)$ can be any given left continuous function in t . μ is a given constant. The data must contain some right censored observations. If there is no censoring or the only censoring is the largest observation, the code will stop and we should use `el.test()` which is for uncensored data.

The log empirical likelihood been maximized is

$$\sum_{d_i=1} \log \Delta F(x_i) + \sum_{d_i=0} \log [1 - F(x_i)].$$

Usage

```
el.cen.test(x,d,fun=function(x){x},mu,error=1e-8,maxit=15)
```

Arguments

x	a vector containing the observed survival times.
d	a vector containing the censoring indicators, 1-uncensor; 0-censor.
fun	a left continuous (weight) function used to calculate the mean as in H_0 . <code>fun(t)</code> must be able to take a vector input <code>t</code> . Default to the identity function $f(t) = t$.
mu	a real number used in the constraint, sum to this value.
error	an optional positive real number specifying the tolerance of iteration error in the QP. This is the bound of the L_1 norm of the difference of two successive weights.
maxit	an optional integer, used to control maximum number of iterations.

Details

When the given constants μ is too far away from the NPMLE, there will be no distribution satisfy the constraint. In this case the computation will stop. The $-2 \log$ empirical likelihood ratio should be infinite.

The constant `mu` must be inside $(\min f(x_i), \max f(x_i))$ for the computation to continue. It is always true that the NPMLE values are feasible. So when the computation cannot continue, try move the `mu` closer to the NPMLE, or use a different `fun`.

This function depends on `Wdataclean2()`, `WKM()` and `solve3.QP()`

This function uses sequential Quadratic Programming to find the maximum. Unlike other functions in this package, it can be slow for larger sample sizes. It took about one minute for a sample of size 2000 with 20% censoring on a 1GHz, 256MB PC, about 19 seconds on a 3 GHz 512MB PC.

Value

A list with the following components:

"-2LLR"	The -2Log Likelihood ratio.
xtimes	the location of the CDF jumps.
weights	the jump size of CDF at those locations.
Pval	P-value
error	the L_1 norm between the last two wts.
iteration	number of iterations carried out

Author(s)

Mai Zhou, Kun Chen

References

Pan, X. and Zhou, M. (1999). Using 1-parameter sub-family of distributions in empirical likelihood ratio with censored data. *J. Statist. Plann. Inference*, **75**, 379-392.

Chen, K. and Zhou, M. (2000). Computing censored empirical likelihood ratio using Quadratic Programming. *Tech Report, Univ. of Kentucky, Dept of Statistics*

Zhou, M. and Chen, K. (2007). Computation of the empirical likelihood ratio from censored data. *Journal of Statistical Computing and Simulation*, **77**, 1033-1042.

Examples

```
e1.cen.test(rexp(100), c(rep(0,25),rep(1,75)), mu=1.5)
## second example with tied observations
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
e1.cen.test(x,d,mu=3.5)
# we should get "-2LLR" = 1.246634 etc.
```

 el.ltrc.EM

Empirical likelihood ratio for mean with left truncated and right censored data, by EM algorithm

Description

This program uses EM algorithm to compute the maximized (wrt p_i) empirical log likelihood function for left truncated and right censored data with the MEAN constraint:

$$\sum_{d_i=1} p_i f(x_i) = \int f(t) dF(t) = \mu .$$

Where $p_i = \Delta F(x_i)$ is a probability, d_i is the censoring indicator, 1(uncensored), 0(right censored). The d for the largest observation x , is always (automatically) changed to 1. μ is a given constant. This function also returns those p_i .

The log empirical likelihood function been maximized is

$$\sum_{d_i=1} \log \frac{\Delta F(x_i)}{1 - F(y_i)} + \sum_{d_i=0} \log \frac{1 - F(x_i)}{1 - F(y_i)} .$$

Usage

```
el.ltrc.EM(y,x,d,fun=function(t){t},mu,maxit=30,error=1e-9)
```

Arguments

y	an optional vector containing the observed left truncation times.
x	a vector containing the censored survival times.
d	a vector containing the censoring indicators, 1-uncensored; 0-right censored.
fun	a continuous (weight) function used to calculate the mean as in H_0 . fun(t) must be able to take a vector input t. Default to the identity function $f(t) = t$.
mu	a real number used in the constraint, mean value of $f(X)$.
error	an optional positive real number specifying the tolerance of iteration error. This is the bound of the L_1 norm of the difference of two successive weights.
maxit	an optional integer, used to control maximum number of iterations.

Details

We return the -2 log likelihood ratio, and the constrained NPMLE of CDF. The un-constrained NPMLE should be WJT or Lynden-Bell estimator.

When the given constants μ is too far away from the NPMLE, there will be no distribution satisfy the constraint. In this case the computation will stop. The -2 Log empirical likelihood ratio should be infinite.

The constant mu must be inside $(\min f(x_i), \max f(x_i))$ for the computation to continue. It is always true that the NPMLE values are feasible. So when the computation stops, try move the mu closer to the NPMLE —

$$\sum_{d_i=1} p_i^0 f(x_i)$$

p_i^0 taken to be the jumps of the NPMLE of CDF. Or use a different fun.

This implementation is all in R and have several for-loops in it. A faster version would use C to do the for-loop part. (but this version is easier to port to Splus, and seems faster enough).

Value

A list with the following components:

times	locations of CDF that have positive mass.
prob	the probability of the constrained NPMLE of CDF at those locations.
"-2LLR"	It is Minus two times the Empirical Log Likelihood Ratio. Should be approximate chi-square distributed under Ho.

Author(s)

Mai Zhou

References

- Zhou, M. (2002). Computing censored and truncated empirical likelihood ratio by EM algorithm. *Tech Report, Univ. of Kentucky, Dept of Statistics*
- Tsai, W. Y., Jewell, N. P., and Wang, M. C. (1987). A note on product-limit estimator under right censoring and left truncation. *Biometrika*, **74**, 883-886.
- Turnbull, B. (1976). The empirical distribution function with arbitrarily grouped, censored and truncated data. *JRSS B*, 290-295.
- Zhou, M. (2005). Empirical likelihood ratio with arbitrarily censored/truncated data by EM algorithm. *Journal of Computational and Graphical Statistics* **14**, 643-656.

Examples

```
## example with tied observations
y <- c(0, 0, 0.5, 0, 1, 2, 2, 0, 0, 0, 0, 0 )
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
e1.ltrc.EM(y,x,d,mu=3.5)
ypsy <- c(51, 58, 55, 28, 25, 48, 47, 25, 31, 30, 33, 43, 45, 35, 36)
xpsy <- c(52, 59, 57, 50, 57, 59, 61, 61, 62, 67, 68, 69, 69, 65, 76)
dpsy <- c(1, 1, 1, 1, 1, 1, 1, 1, 0, 0, 0, 1, 1, 0, 1 )
e1.ltrc.EM(ypsy,xpsy,dpsy,mu=64)
```

 el.test

Empirical likelihood ratio test for the means, uncensored data

Description

Compute the empirical likelihood ratio with the mean vector fixed at mu.

The log empirical likelihood been maximized is

$$\sum_{i=1}^n \log \Delta F(x_i).$$

Usage

```
el.test(x, mu, lam, maxit=25, gradtol=1e-7,
        svdtol = 1e-9, itertrace=FALSE)
```

Arguments

x	a matrix or vector containing the data, one row per observation.
mu	a numeric vector (of length = ncol(x)) to be tested as the mean vector of x above, as H_0 .
lam	an optional vector of length = length(mu), the starting value of Lagrange multipliers, will use 0 if missing.
maxit	an optional integer to control iteration when solve constrained maximization.
gradtol	an optional real value for convergence test.
svdtol	an optional real value to detect singularity while solve equations.
itertrace	a logical value. If the iteration history needs to be printed out.

Details

If mu is in the interior of the convex hull of the observations x, then wts should sum to n. If mu is outside the convex hull then wts should sum to nearly zero, and -2LLR will be a large positive number. It should be infinity, but for inferential purposes a very large number is essentially equivalent. If mu is on the boundary of the convex hull then wts should sum to nearly k where k is the number of observations within that face of the convex hull which contains mu.

When mu is interior to the convex hull, it is typical for the algorithm to converge quadratically to the solution, perhaps after a few iterations of searching to get near the solution. When mu is outside or near the boundary of the convex hull, then the solution involves a lambda of infinite norm. The algorithm tends to nearly double lambda at each iteration and the gradient size then decreases roughly by half at each iteration.

The goal in writing the algorithm was to have it “fail gracefully” when mu is not inside the convex hull. The user can either leave -2LLR “large and positive” or can replace it by infinity when the weights do not sum to nearly n.

Value

A list with the following components:

-2LLR	the -2 loglikelihood ratio; approximate chisq distribution under H_o .
Pval	the observed P-value by chi-square approximation.
lambda	the final value of Lagrange multiplier.
grad	the gradient at the maximum.
hess	the Hessian matrix.
wt	weights on the observations
nits	number of iteration performed

Author(s)

Original Splus code by Art Owen. Adapted to R by Mai Zhou.

References

Owen, A. (1990). Empirical likelihood ratio confidence regions. *Ann. Statist.* **18**, 90-120.

Examples

```
x <- matrix(c(rnorm(50,mean=1), rnorm(50,mean=2)), ncol=2,nrow=50)
el.test(x, mu=c(1,2))
## Suppose now we wish to test Ho: 2mu(1)-mu(2)=0, then
y <- 2*x[,1]-x[,2]
el.test(y, mu=0)
xx <- c(28,-44,29,30,26,27,22,23,33,16,24,29,24,40,21,31,34,-2,25,19)
el.test(xx, mu=15) ##### -2LLR = 1.805702
```

el.test.wt

Weighted Empirical Likelihood ratio for mean, uncensored data

Description

This program is similar to `el.test()` except it takes weights, and is for one dimensional mu.

The mean constraint considered is:

$$\sum_{i=1}^n p_i x_i = \mu.$$

where $p_i = \Delta F(x_i)$ is a probability. Plus the probability constraint: $\sum p_i = 1$.

The weighted log empirical likelihood been maximized is

$$\sum_{i=1}^n w_i \log p_i.$$

Usage

```
el.test.wt(x, wt, mu, usingC=TRUE)
```

Arguments

x	a vector containing the observations.
wt	a vector containing the weights.
mu	a real number used in the constraint, weighted mean value of $f(X)$.
usingC	TRUE: use C function, which may be benefit when sample size is large; FALSE: use pure R function.

Details

This function used to be an internal function. It becomes external because others may find it useful elsewhere.

The constant mu must be inside $(\min x_i, \max x_i)$ for the computation to continue.

Value

A list with the following components:

x	the observations.
wt	the vector of weights.
prob	The probabilities that maximized the weighted empirical likelihood under mean constraint.

Author(s)

Mai Zhou, Y.F. Yang for C part.

References

Owen, A. (1990). Empirical likelihood ratio confidence regions. *Ann. Statist.* **18**, 90-120.

Zhou, M. (2002). Computing censored empirical likelihood ratio by EM algorithm. *Tech Report, Univ. of Kentucky, Dept of Statistics*

Examples

```
## example with tied observations
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
el.cen.EM(x,d,mu=3.5)
## we should get "-2LLR" = 1.2466...
myfun5 <- function(x, theta, eps) {
  u <- (x-theta)*sqrt(5)/eps
  INDE <- (u < sqrt(5)) & (u > -sqrt(5))
  u[u >= sqrt(5)] <- 0
  u[u <= -sqrt(5)] <- 1
}
```



```

y <- 0.5 - (u - (u)^3/15)*3/(4*sqrt(5))
u[ INDE ] <- y[ INDE ]
return(u)
}
el.cen.EM(x, d, fun=myfun5, mu=0.5, theta=3.5, eps=0.1)

```

el.test.wt2

*Weighted Empirical Likelihood ratio for mean(s), uncensored data***Description**

This program is similar to `el.test()` except it takes weights.

The mean constraints are:

$$\sum_{i=1}^n p_i x_i = \mu.$$

Where $p_i = \Delta F(x_i)$ is a probability. Plus the probability constraint: $\sum p_i = 1$.

The weighted log empirical likelihood been maximized is

$$\sum_{i=1}^n w_i \log p_i.$$

Usage

```

el.test.wt2(x, wt, mu, maxit = 25, gradtol = 1e-07, Hessian = FALSE,
            svdtol = 1e-09, itertrace = FALSE)

```

Arguments

<code>x</code>	a matrix (of size $n \times p$) or vector containing the observations.
<code>wt</code>	a vector of length n , containing the weights. If weights are all 1, this is very similar to <code>el.test</code> . <code>wt</code> have to be positive.
<code>mu</code>	a vector of length p , used in the constraint. weighted mean value of $f(X)$.
<code>maxit</code>	an integer, the maximum number of iteration.
<code>gradtol</code>	a positive real number, the tolerance for a solution
<code>Hessian</code>	logical. if the Hessian needs to be computed?
<code>svdtol</code>	tolerance in perform SVD of the Hessian matrix.
<code>itertrace</code>	TRUE/FALSE, if the intermediate steps needs to be printed.

Details

This function used to be an internal function. It becomes external because others may find it useful.

It is similar to the function `el.test()` with the following differences:

- (1) The output `lambda` in `el.test.wts`, when divided by n (the sample size or sum of all the weights) should be equal to the output `lambda` in `el.test`.
- (2) The Newton step of iteration in `el.test.wts` is different from those in `el.test`. (even when all the weights are one).

Value

A list with the following components:

lambda	the Lagrange multiplier. Solution.
wt	the vector of weights.
grad	The gradian at the final solution.
nits	number of iterations performed.
prob	The probabilities that maximized the weighted empirical likelihood under mean constraint.

Author(s)

Mai Zhou

References

- Owen, A. (1990). Empirical likelihood ratio confidence regions. *Ann. Statist.* **18**, 90-120.
- Zhou, M. (2005). Empirical likelihood ratio with arbitrary censored/truncated data by EM algorithm. *Journal of Computational and Graphical Statistics*, **14**, 643-656.
- Zhou, M. (2002). Computing censored empirical likelihood ratio by EM algorithm. *Tech Report, Univ. of Kentucky, Dept of Statistics*

Examples

```
## example with tied observations
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
el.cen.EM(x,d,mu=3.5)
## we should get "-2LLR" = 1.2466...
myfun5 <- function(x, theta, eps) {
  u <- (x-theta)*sqrt(5)/eps
  INDE <- (u < sqrt(5)) & (u > -sqrt(5))
  u[u >= sqrt(5)] <- 0
  u[u <= -sqrt(5)] <- 1
  y <- 0.5 - (u - (u)^3/15)*3/(4*sqrt(5))
  u[ INDE ] <- y[ INDE ]
  return(u)
}
el.cen.EM(x, d, fun=myfun5, mu=0.5, theta=3.5, eps=0.1)
```

el.trun.test *Empirical likelihood ratio for mean with left truncated data*

Description

This program uses EM algorithm to compute the maximized (wrt p_i) empirical log likelihood function for left truncated data with the MEAN constraint:

$$\sum p_i f(x_i) = \int f(t) dF(t) = \mu .$$

Where $p_i = \Delta F(x_i)$ is a probability. μ is a given constant. It also returns those p_i and the p_i without constraint, the Lynden-Bell estimator.

The log likelihood been maximized is

$$\sum_{i=1}^n \log \frac{\Delta F(x_i)}{1 - F(y_i)} .$$

Usage

```
el.trun.test(y,x,fun=function(t){t},mu,maxit=20,error=1e-9)
```

Arguments

y	a vector containing the left truncation times.
x	a vector containing the survival times. truncation means $x > y$.
fun	a continuous (weight) function used to calculate the mean as in H_0 . $\text{fun}(t)$ must be able to take a vector input t . Default to the identity function $f(t) = t$.
mu	a real number used in the constraint, mean value of $f(X)$.
error	an optional positive real number specifying the tolerance of iteration error. This is the bound of the L_1 norm of the difference of two successive weights.
maxit	an optional integer, used to control maximum number of iterations.

Details

This implementation is all in R and have several for-loops in it. A faster version would use C to do the for-loop part. But it seems faster enough and is easier to port to Splus.

When the given constants μ is too far away from the NPMLE, there will be no distribution satisfy the constraint. In this case the computation will stop. The -2 Log empirical likelihood ratio should be infinite.

The constant mu must be inside $(\min f(x_i), \max f(x_i))$ for the computation to continue. It is always true that the NPMLE values are feasible. So when the computation stops, try move the mu closer to the NPMLE —

$$\sum_{d_i=1}^0 p_i^0 f(x_i)$$

p_i^0 taken to be the jumps of the NPMLE of CDF. Or use a different fun.

Usage

```
emplikH.disc(x, d, y= -Inf, K, fun, tola=.Machine$double.eps^.25, theta)
```

Arguments

x a vector, the observed survival times.

d a vector, the censoring indicators, 1-uncensor; 0-censor.

y optional vector, the left truncation times.

K a real number used in the constraint, sum to this value.

fun a left continuous (weight) function used to calculate the weighted discrete hazard in H_0 . `fun(x, theta)` must be able to take a vector input `x`, and a parameter `theta`.

tola an optional positive real number specifying the tolerance of iteration error in solve the non-linear equation needed in constrained maximization.

theta a given real number used as the parameter of the function f .

Details

The log likelihood been maximized is the ‘binomial’ empirical likelihood:

$$\sum D_i \log w_i + (R_i - D_i) \log[1 - w_i]$$

where $w_i = \Delta H(t_i)$ is the jump of the cumulative hazard function, D_i is the number of failures observed at t_i , R_i is the number of subjects at risk at time t_i .

For discrete distributions, the jump size of the cumulative hazard at the last jump is always 1. We have to exclude this jump from the summation since $\log(1 - dH(\cdot))$ do not make sense.

The constants `theta` and `K` must be inside the so called feasible region for the computation to continue. This is similar to the requirement that in testing the value of the mean, the value must be inside the convex hull of the observations. It is always true that the NPMLE values are feasible. So when the computation stops, try move the `theta` and `K` closer to the NPMLE. When the computation stops, the $-2LLR$ should have value infinite.

In case you do not need the `theta` in the definition of the function f , you still need to formally define your `fun` function with a `theta` input, just to match the arguments.

Value

A list with the following components:

times the location of the hazard jumps.

wts the jump size of hazard function at those locations.

lambda the final value of the Lagrange multiplier.

"-2LLR" The discrete -2Log Likelihood ratio.

Pval P-value

niters number of iterations used

Author(s)

Mai Zhou

References

Fang, H. (2000). Binomial Empirical Likelihood Ratio Method in Survival Analysis. Ph.D. Thesis, Univ. of Kentucky, Dept of Statistics.

Zhou and Fang (2001). "Empirical likelihood ratio for 2 sample problem for censored data". *Tech Report, Univ. of Kentucky, Dept of Statistics*

Zhou, M. and Fang, H. (2006). A comparison of Poisson and binomial empirical likelihood. *Tech Report, Univ. of Kentucky, Dept of Statistics*

Examples

```
fun4 <- function(x, theta) { as.numeric(x <= theta) }
x <- c(1, 2, 3, 4, 5, 6, 5, 4, 3, 4, 1, 2.4, 4.5)
d <- c(1, 0, 1, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1)
# test if -H(4) = -0.7
emplikH.disc(x=x,d=d,K=-0.7,fun=fun4,theta=4)
# we should get "-2LLR" 0.1446316 etc...
y <- c(-2,-2, -2, 1.5, -1)
emplikH.disc(x=x,d=d,y=y,K=-0.7,fun=fun4,theta=4)
```

 emplikH.disc2

Two sample empirical likelihood ratio for discrete hazards with right censored, left truncated data, one parameter.

Description

Use empirical likelihood ratio and Wilks theorem to test the null hypothesis that

$$\int f_1(t)I_{[dH_1 < 1]} \log(1 - dH_1(t)) - \int f_2(t)I_{[dH_2 < 1]} \log(1 - dH_2(t)) = \theta$$

where $H_*(t)$ is the (unknown) discrete cumulative hazard function; $f_*(t)$ can be any predictable functions of t . θ is the parameter. The given value of θ in these computation is the value to be tested. The data can be right censored and left truncated.

When the given constants θ is too far away from the NPMLE, there will be no hazard function satisfy this constraint and the -2 Log empirical likelihood ratio will be infinite. In this case the computation will stop.

Usage

```
emplikH.disc2(x1, d1, y1= -Inf, x2, d2, y2 = -Inf,
              theta, fun1, fun2, tola = 1e-6, maxi, mini)
```

Arguments

x1	a vector, the observed survival times, sample 1.
d1	a vector, the censoring indicators, 1-uncensor; 0-censor.
y1	optional vector, the left truncation times.
x2	a vector, the observed survival times, sample 2.
d2	a vector, the censoring indicators, 1-uncensor; 0-censor.
y2	optional vector, the left truncation times.
fun1	a predictable function used to calculate the weighted discrete hazard in H_0 . fun1(x) must be able to take a vector input x.
fun2	similar to fun1, but for sample 2.
tola	an optional positive real number, the tolerance of iteration error in solve the non-linear equation needed in constrained maximization.
theta	a given real number. for Ho constraint.
maxi	upper bound for lambda, usually positive.
mini	lower bound for lambda, usually negative.

Details

The log likelihood been maximized is the ‘binomial’ empirical likelihood:

$$\sum D_{1i} \log w_i + (R_{1i} - D_{1i}) \log[1 - w_i] + \sum D_{2j} \log v_j + (R_{2j} - D_{2j}) \log[1 - v_j]$$

where $w_i = \Delta H_1(t_i)$ is the jump of the cumulative hazard function at t_i , D_{1i} is the number of failures observed at t_i , R_{1i} is the number of subjects at risk at time t_i .

For discrete distributions, the jump size of the cumulative hazard at the last jump is always 1. We have to exclude this jump from the summation in the constraint calculation since $\log(1 - dH(\cdot))$ do not make sense.

The constants theta must be inside the so called feasible region for the computation to continue. This is similar to the requirement that in ELR testing the value of the mean, the value must be inside the convex hull of the observations. It is always true that the NPMLE values are feasible. So when the computation stops, try move the theta closer to the NPMLE. When the computation stops, the -2LLR should have value infinite.

Value

A list with the following components:

times	the location of the hazard jumps.
wts	the jump size of hazard function at those locations.
lambda	the final value of the Lagrange multiplier.
"-2LLR"	The -2Log Likelihood ratio.
Pval	P-value
niters	number of iterations used

Author(s)

Mai Zhou

References

Zhou and Fang (2001). "Empirical likelihood ratio for 2 sample problems for censored data". *Tech Report, Univ. of Kentucky, Dept of Statistics*

Examples

```

if(require("boot", quietly = TRUE)) {
###library(boot)
data(channing)
ymale <- channing[1:97,2]
dmale <- channing[1:97,5]
xmale <- channing[1:97,3]
yfemal <- channing[98:462,2]
dfemal <- channing[98:462,5]
xfemal <- channing[98:462,3]
fun1 <- function(x) { as.numeric(x <= 960) }
emplikH.disc2(x1=xfemal, d1=dfemal, y1=yfemal,
  x2=xmale, d2=dmale, y2=ymale, theta=0.2, fun1=fun1, fun2=fun1, maxi=4, mini=-10)
#####
### You should get "-2LLR" = 1.511239 and a lot more other outputs.
#####
emplikH.disc2(x1=xfemal, d1=dfemal, y1=yfemal,
  x2=xmale, d2=dmale, y2=ymale, theta=0.25, fun1=fun1, fun2=fun1, maxi=4, mini=-5)
#####
### This time you get "-2LLR" = 1.150098 etc. etc.
#####
}

```

emplikH1.test

Empirical likelihood for hazard with right censored, left truncated data

Description

Use empirical likelihood ratio and Wilks theorem to test the null hypothesis that

$$\int f(t)dH(t) = \theta$$

with right censored, left truncated data. Where $H(t)$ is the unknown cumulative hazard function; $f(t)$ can be any given function and θ a given constant. In fact, $f(t)$ can even be data dependent, just have to be 'predictable'.

Usage

```
emplikH1.test(x, d, y= -Inf, theta, fun, tola=.Machine$double.eps^.5)
```


Arguments

x	a vector of the censored survival times.
d	a vector of the censoring indicators, 1-uncensor; 0-censor.
y	a vector of the observed left truncation times.
theta	a real number used in the H_0 to set the hazard to this value.
fun	a left continuous (weight) function used to calculate the weighted hazard in H_0 . fun must be able to take a vector input. See example below.
tola	an optional positive real number specifying the tolerance of iteration error in solve the non-linear equation needed in constrained maximization.

Details

This function is designed for the case where the true distributions are all continuous. So there should be no tie in the data.

The log empirical likelihood used here is the ‘Poisson’ version empirical likelihood:

$$\sum_{i=1}^n \delta_i \log(dH(x_i)) - [H(x_i) - H(y_i)] .$$

If there are ties in the data that are resulted from rounding, you may break the tie by adding a different tiny number to the tied observation(s). If those are true ties (thus the true distribution is discrete) we recommend use `emplikdisc.test()`.

The constant theta must be inside the so called feasible region for the computation to continue. This is similar to the requirement that in testing the value of the mean, the value must be inside the convex hull of the observations. It is always true that the NPMLE values are feasible. So when the computation complains that there is no hazard function satisfy the constraint, you should try to move the theta value closer to the NPMLE. When the computation stops prematurely, the -2LLR should have value infinite.

Value

A list with the following components:

times	the location of the hazard jumps.
wts	the jump size of hazard function at those locations.
lambda	the Lagrange multiplier.
"-2LLR"	the -2Log Likelihood ratio.
Pval	P-value
niters	number of iterations used

Author(s)

Mai Zhou

References

Pan, X. and Zhou, M. (2002), “Empirical likelihood in terms of hazard for censored data”. *Journal of Multivariate Analysis* **80**, 166-188.

Examples

```
fun <- function(x) { as.numeric(x <= 6.5) }
emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=2, fun=fun)
fun2 <- function(x) {exp(-x)}
emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=0.2, fun=fun2)
```

emplikH1B	<i>Return binomial empirical likelihood ratio for the given lambda, with right censored data</i>
-----------	--

Description

Compute the binomial empirical likelihood ratio for the given tilt parameter lambda. Most useful for construct Wilks confidence intervals. The null hypothesis or constraint is defined by the parameter θ , where

$$\int fung(t) d\log(1 - H(t)) = \theta$$

Where $H(t)$ is the unknown cumulative hazard function; $fung(t)$ can be any given function. In the future, the function $fung$ may be replaced by the vector of $fung(x)$, since this is more flexible.

Input data can be right censored. If no censoring, set $d=\text{rep}(1, \text{length}(x))$.

Usage

```
emplikH1B(lambda, x, d, fung, CIforTheta=FALSE)
```

Arguments

lambda	a scalar. Can be positive or negative. The amount of tiling.
x	a vector of the censored survival times.
d	a vector of the censoring indicators, 1-uncensor; 0-right censor.
fung	a left continuous (weight) function used to calculate the weighted hazard in the parameter θ . fung must be able to take a vector input. See example below.
CIforTheta	an optional logical value. Default to FALSE. If set to TRUE, will return the integrated hazard value for the given lambda.

Details

This function is used to calculate lambda confidence interval (Wilks type) for θ .

This function is designed for the case where the true distribution should be discrete. Ties in the data are OK.

The log empirical likelihood used here is the ‘binomial’ version empirical likelihood:

$$\sum_{i=1}^n \delta_i \log(dH(x_i)) + (R_i - \delta_i) \log[1 - dH(x_i)].$$

Value

A list with the following components:

times	the location of the hazard jumps.
jumps	the jump size of hazard function at those locations.
lambda	the input lambda.
"-2LLR"	the -2Log Likelihood ratio.
IntHaz	The theta defined above, for the given lambda.

Author(s)

Mai Zhou

References

Pan, X. and Zhou, M. (2002), “Empirical likelihood in terms of hazard for censored data”. *Journal of Multivariate Analysis* **80**, 166-188.

Examples

```
## fun <- function(x) { as.numeric(x <= 6.5) }
## emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=2, fun=fun)
## fun2 <- function(x) {exp(-x)}
## emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=0.2, fun=fun2)
```

emplikHIP

Return Poisson Empirical likelihood ratio for the given lambda, with right censored data

Description

Compute the Poisson empirical likelihood ratio for the given tilt parameter lambda. Most useful for the construction of Wilks confidence intervals. The null hypothesis or constraint is defined by the parameter θ , where

$$\int \text{fung}(t)dH(t) = \theta$$

Where $H(t)$ is the unknown cumulative hazard function; $\text{fung}(t)$ can be any given function.

In the future, the function fung may be replaced by the vector of $\text{fung}(x)$, since this is more flexible.

Input data can be right censored. If no censoring, set $d=\text{rep}(1, \text{length}(x))$.

Usage

```
emplikH1P(lambda, x, d, fung, CIforTheta=FALSE)
```

Arguments

lambda	a scalar. Can be positive or negative. The amount of tiling.
x	a vector of the censored survival times.
d	a vector of the censoring indicators, 1-uncensor; 0-right censor.
fung	a left continuous (weight) function used to calculate the weighted hazard in the parameter θ . fung must be able to take a vector input. See example below.
CIforTheta	an optional logical value. Default to FALSE. If set to TRUE, will return the integrated hazard value for the given lambda.

Details

This function is for calculate lambda confidence intervals for θ .

This function is designed for the case where the true distribution should be continuous. So there should be no tie in the data.

The log empirical likelihood used here is the ‘Poisson’ version empirical likelihood:

$$\sum_{i=1}^n \delta_i \log(dH(x_i)) - [H(x_i)].$$

If there are ties in the data that are resulted from rounding, you may want to break the tie by adding a different tiny number to the tied observation(s). For example: 2, 2, 2, change to 2.00001, 2.00002, 2.00003. If those are true ties (thus the true distribution must be discrete) we recommend to use `emplikH1B` instead.

Value

A list with the following components:

times	the location of the hazard jumps.
wts	the jump size of hazard function at those locations.

lambda the Lagrange multiplier.
 "-2LLR" the -2Log Empirical Likelihood ratio, Poisson version.
 MeanHaz The theta defined above, the hazard integral, if CIforTheta =TRUE.

Author(s)

Mai Zhou

References

Pan, X. and Zhou, M. (2002), "Empirical likelihood in terms of hazard for censored data". *Journal of Multivariate Analysis* **80**, 166-188.

Examples

```
## fun <- function(x) { as.numeric(x <= 6.5) }
## emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=2, fun=fun)
## fun2 <- function(x) {exp(-x)}
## emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=0.2, fun=fun2)
```

emplikH2.test	<i>Empirical likelihood for weighted hazard with right censored, left truncated data</i>
---------------	--

Description

Use empirical likelihood ratio and Wilks theorem to test the null hypothesis that

$$\int f(t, \dots) dH(t) = K$$

with right censored, left truncated data, where $H(t)$ is the (unknown) cumulative hazard function; $f(t, \dots)$ can be any given left continuous function in t ; (of course the integral must be finite).

Usage

```
emplikH2.test(x, d, y= -Inf, K, fun, tola=.Machine$double.eps^.5, ...)
```

Arguments

x a vector containing the censored survival times.
 d a vector of the censoring indicators, 1-uncensor; 0-censor.
 y a vector containing the left truncation times. If left as default value, -Inf, it means no truncation.
 K a real number used in the constraint, i.e. to set the weighted integral of hazard to this value.

fun	a left continuous (in t) weight function used to calculate the weighted hazard in H_0 . $\text{fun}(t, \dots)$ must be able to take a vector input t .
tola	an optional positive real number specifying the tolerance of iteration error in solve the non-linear equation needed in constrained maximization.
...	additional parameter(s), if any, passing along to fun. This allows an implicit function of fun.

Details

This version works for implicit function $f(t, \dots)$.

This function is designed for continuous distributions. Thus we do not expect tie in the observation x . If you believe the true underlying distribution is continuous but the sample observations have tie due to rounding, then you might want to add a small number to the observations to break tie.

The likelihood used here is the ‘Poisson’ version of the empirical likelihood

$$\prod_{i=1}^n (dH(x_i))^{\delta_i} \exp[-H(x_i) + H(y_i)].$$

For discrete distributions we recommend use `emplikdisc.test()`.

Please note here the largest observed time is NOT automatically defined to be uncensored. In the `el.cen.EM()`, it is (to make F a proper distribution always).

The constant K must be inside the so called feasible region for the computation to continue. This is similar to the requirement that when testing the value of the mean, the value must be inside the convex hull of the observations for the computation to continue. It is always true that the NPMLE value is feasible. So when the computation cannot continue, that means there is no hazard function dominated by the Nelson-Aalen estimator satisfy the constraint. You may try to move the θ and K closer to the NPMLE. When the computation cannot continue, the $-2LLR$ should have value infinite.

Value

A list with the following components:

times	the location of the hazard jumps.
wts	the jump size of hazard function at those locations.
lambda	the Lagrange multiplier.
"-2LLR"	the $-2\text{Log Likelihood ratio}$.
Pval	P-value
niters	number of iterations used

Author(s)

Mai Zhou

References

Pan, XR and Zhou, M. (2002), “Empirical likelihood in terms of cumulative hazard for censored data”. *Journal of Multivariate Analysis* **80**, 166-188.

See Also

emplikHs.test2

Examples

```
z1<-c(1,2,3,4,5)
d1<-c(1,1,0,1,1)
fun4 <- function(x, theta) { as.numeric(x <= theta) }
emplikH2.test(x=z1,d=d1, K=0.5, fun=fun4, theta=3.5)
#Next, test if H(3.5) = log(2) .
emplikH2.test(x=z1,d=d1, K=log(2), fun=fun4, theta=3.5)
#Next, try one sample log rank test
indi <- function(x,y){ as.numeric(x >= y) }
fun3 <- function(t,z){rowsum(outer(z,t,FUN="indi"),group=rep(1,length(z)))}
emplikH2.test(x=z1, d=d1, K=sum(0.25*z1), fun=fun3, z=z1)
##this is testing if the data is from an exp(0.25) population.
```

emplikH2B

Return binomial empirical likelihood ratio for the given lambda, with 2-sample right censored data

Description

Compute the binomial empirical likelihood ratio for the given tilt parameter lambda. Most useful for construct Wilks confidence intervals. The null hypothesis or constraint is defined by the parameter θ , where

$$\int fun1(t)d\log(1 - H_1(t)) - \int fun2(t)d\log(1 - H_2(t)) = \theta$$

If the lambda=0, you get the Nelson-Aalen (NPMLE) and output -2LLR =0. Otherwise the lambda is not scaled (as in one sample case). Since there are two sample sizes. It can be confusing which sample size to use for scale. So the lambda here is larger than those in one sample by a sclae of (either?) sample size.

Where $H_1(t)$ and $H_2(t)$ are the unknown cumulative hazard function for sample 1/2; $fun1(t)$ and $fun2(t)$ can be any given function. It can even be random, just need to be predictable. In the future, the input function fun may replaced by the vector of $fun(x)$, since this is more flexible.

Input data can be right censored. If no censoring, set $d1=rep(1, length(x1))$, and/or $d2=rep(1, length(x2))$.

Usage

```
emplikH2B(lambda, x1, d1, x2, d2, fun1, fun2, CIforTheta=FALSE)
```

Arguments

lambda	a scalar. Can be positive or negative. The amount of tiling.
x1	a vector of the censored survival times. sample 1
d1	a vector of the censoring indicators, 1-uncensor; 0-right censor.
x2	a vector of the censored survival times. sample 2
d2	a vector of the censoring indicators, 1-uncensor; 0-right censor.
fun1	a left continuous (weight) function used to calculate the weighted hazard in the parameter θ . fun1 must be able to take a vector input. See example below.
fun2	Ditto
CIforTheta	an optional logical value. Default to FALSE. If set to TRUE, will return the integrated hazard value for the given lambda.

Details

This function is used to calculate lambda confidence interval (Wilks type) for θ .

This function is designed for the case where the true distribution should be discrete. Ties in the data are OK.

The log empirical likelihood used here is the ‘binomial’ version empirical likelihood:

$$\log EL1 = \sum_{i=1}^n \delta_i \log(dH(x_i)) + (R_i - \delta_i) \log[1 - dH(x_i)],$$

(similarly defined for sample 2) and the overall log EL = log EL1 + log EL2.

Value

A list with the following components:

"-2LLR"	the -2Log Empirical Likelihood ratio, binomial version.
lambda	the input lambda. The tilt. The Lagrange multiplier.
times1	the location of the hazard jumps. sample 1.
times2	the location of the hazard jumps. sample 2.
wts1	the jump size of hazard function at those locations.
wts2	the jump size of hazard function at those locations.
HazDiff2	Difference of two hazard integrals. theta defined above.

Author(s)

Mai Zhou

References

Pan, X. and Zhou, M. (2002), “Empirical likelihood in terms of hazard for censored data”. *Journal of Multivariate Analysis* **80**, 166-188.

Examples

```
## fun <- function(x) { as.numeric(x <= 6.5) }
## emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=2, fun=fun)
## fun2 <- function(x) {exp(-x)}
## emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=0.2, fun=fun2)
```

emplikH2P	<i>Return Poisson Empirical likelihood ratio for the given lambda, with 2-sample right censored data</i>
-----------	--

Description

Compute the Poisson empirical likelihood ratio for the given tilt parameter lambda. Most useful when construct Wilks confidence intervals. The null hypothesis or constraint is defined by the parameter θ , where

$$\int fun1(t)dH_1(t) - \int fun2(t)dH_2(t) = \theta$$

If the lambda value set to zero, you get the Nelson-Aalen (NPMLE) and output $-2LLR = 0$. For other values of lambda, this is not scaled (like in one sample) since there are two samples and it can be confusing which sample size to use. So here, the lambda is larger (with a scale of sample size) than those in one sample: emplikH1P.

Where $H_1(t)$ is the unknown cumulative hazard function of sample one; $H_2(t)$ is the cumulative function of sample two; $fun1(t)$ can be any given function. In the future, the function $fun1$ may be replaced by the vector of $fun(x)$, since this is more flexible. Same for $fun2$.

Input data can be right censored. If no censoring, set $d1 = \text{rep}(1, \text{length}(x1))$, and/or $d2 = \text{rep}(1, \text{length}(x2))$.

Usage

```
emplikH2P(lambda, x1, d1, x2, d2, fun1, fun2, CIforTheta=FALSE)
```

Arguments

lambda	a scalar. Can be positive or negative. The amount of tiling.
x1	a vector of the censored survival times. sample one.
d1	a vector of the censoring indicators, 1-uncensor; 0-right censor.
x2	a vector of the censored survival times. sample two.
d2	a vector of the censoring indicators, 1-uncensor; 0-right censor.
fun1	a left continuous (weight) function used to calculate the weighted hazard in the parameter θ . fun1 must be able to take a vector input. See example below.
fun2	Ditto.
CIforTheta	an optional logical value. Default to FALSE. If set to TRUE, will return the integrated hazard value for the given lambda.

Details

This function is for calculate lambda confidence intervals for θ .

This function is designed for the case where the true distribution should be continuous. So there should be no tie in the data.

The log empirical likelihood used here is the ‘Poisson’ version empirical likelihood:

$$EL1 = \sum_{i=1}^n \delta_i \log(dH_1(x_i)) - [H_1(x_i)],$$

(similarly defined for sample 2) and the final EL is the sum of EL1 and EL2.

If there are ties in the data that are resulted from rounding, you may break the tie by adding a different tiny number to the tied observation(s). For example: 2, 2, 2, change to 2.00001, 2.00002, 2.00003. If those are true ties (thus the true distribution must be discrete) we recommend use `emplikH2B`.

Value

A list with the following components:

"-2LLR"	the -2Log Empirical Likelihood ratio, Poisson version.
lambda	The tilt parameter used. It is also the Lagrange multiplier.
"-2LLR(sample1)"	the -2Log EL ratio, sample 1, Poisson version. "-2LLR" = -2LLR(sample1) + -2LLR(sample2)
HazDiff	Average hazard for the constrained hazard integral, if <code>CIforTheta = TRUE</code> .

Author(s)

Mai Zhou

References

Pan, X. and Zhou, M. (2002), “Empirical likelihood in terms of hazard for censored data”. *Journal of Multivariate Analysis* **80**, 166-188.

Examples

```
## fun <- function(x) { as.numeric(x <= 6.5) }
## emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=2, fun=fun)
## fun2 <- function(x) {exp(-x)}
## emplikH1.test( x=c(1,2,3,4,5), d=c(1,1,0,1,1), theta=0.2, fun=fun2)
```

emplikHs.disc2	<i>Two sample empirical likelihood ratio for discrete hazards with right censored, left truncated data. Many constraints.</i>
----------------	---

Description

Use empirical likelihood ratio and Wilks theorem to test the null hypothesis that

$$\int f_1(t)I_{[dH_1 < 1]} \log(1 - dH_1(t)) - \int f_2(t)I_{[dH_2 < 1]} \log(1 - dH_2(t)) = \theta$$

where $H_*(t)$ are the (unknown) discrete cumulative hazard functions; $f_*(t)$ can be any predictable functions of t . θ is a vector of parameters ($\text{dim}=q \geq 1$). The given value of θ in these computation are the value to be tested. The data can be right censored and left truncated.

When the given constants θ is too far away from the NPMLE, there will be no hazard function satisfy this constraint and the -2 Log empirical likelihood ratio will be infinite. In this case the computation will stop.

Usage

```
emplikHs.disc2(x1, d1, y1= -Inf, x2, d2, y2 = -Inf,
              theta, fun1, fun2, maxit=25, tola = 1e-6, itertrace =FALSE)
```

Arguments

x1	a vector, the observed survival times, sample 1.
d1	a vector, the censoring indicators, 1-uncensor; 0-censor.
y1	optional vector, the left truncation times.
x2	a vector, the observed survival times, sample 2.
d2	a vector, the censoring indicators, 1-uncensor; 0-censor.
y2	optional vector, the left truncation times.
fun1	a predictable function used to calculate the weighted discrete hazard in H_0 . fun1(x) must be able to take a vector input (length n) x, and output a matrix of n x q.
fun2	Ditto.
tola	an optional positive real number, the tolerance of iteration error in solve the non-linear equation needed in constrained maximization.
theta	a given vector of length q. for Ho constraint.
maxit	integer, maximum number of iteration.
itertrace	Logocal, lower bound for lambda

Details

The log empirical likelihood been maximized is the ‘binomial empirical likelihood’:

$$\sum D_{1i} \log w_i + (R_{1i} - D_{1i}) \log[1 - w_i] + \sum D_{2j} \log v_j + (R_{2j} - D_{2j}) \log[1 - v_j]$$

where $w_i = \Delta H_1(t_i)$ is the jump of the cumulative hazard function at t_i , D_{1i} is the number of failures observed at t_i , and R_{1i} is the number of subjects at risk at time t_i (for sample one). Similar for sample two.

For discrete distributions, the jump size of the cumulative hazard at the last jump is always 1. We have to exclude this jump from the summation in the constraint calculation since $\log(1 - dH(\cdot))$ do not make sense. In the likelihood, this term contribute a zero (0*Inf).

This function can handle multiple constraints. So $\dim(\theta) = q$. The constants θ must be inside the so called feasible region for the computation to continue. This is similar to the requirement that in testing the value of the mean, the value must be inside the convex hull of the observations. It is always true that the NPMLE values are feasible. So when the computation stops, try move the θ closer to the NPMLE. When the computation stops, the -2LLR should have value infinite.

This code can also be used to compute one sample problems. You need to artificially supply data for sample two (with minimal sample size $(2q+2)$), and supply a function `fun2` that ALWAYS returns zero (zero vector or zero matrix). In the output, read the `-2LLR(sample1)`.

Value

A list with the following components:

<code>times1</code>	the location of the hazard jumps in sample 1.
<code>times2</code>	the location of the hazard jumps in sample 2.
<code>lambda</code>	the final value of the Lagrange multiplier.
<code>"-2LLR"</code>	The -2Log Likelihood ratio.
<code>"-2LLR(sample1)"</code>	The -2Log Likelihood ratio for sample 1 only.
<code>niters</code>	number of iterations used

Author(s)

Mai Zhou

References

Zhou and Fang (2001). “Empirical likelihood ratio for 2 sample problems for censored data”. *Tech Report, Univ. of Kentucky, Dept of Statistics*

Examples

```
if(require("boot", quietly = TRUE)) {
  ###library(boot)
  data(channing)
  ymale <- channing[1:97,2]
```

```

dmale <- channing[1:97,5]
xmale <- channing[1:97,3]
yfemal <- channing[98:462,2]
dfemal <- channing[98:462,5]
xfemal <- channing[98:462,3]
fun1 <- function(x) { as.numeric(x <= 960) }
#####
emplikHs.disc2(x1=xfemal, d1=dfemal, y1=yfemal,
  x2=xmale, d2=dmale, y2=ymale, theta=0.25, fun1=fun1, fun2=fun1)
#####
### This time you get "-2LLR" = 1.150098 etc. etc.
#####
fun2 <- function(x){ cbind(as.numeric(x <= 960), as.numeric(x <= 860))}
##### fun2 has matrix output #####
emplikHs.disc2(x1=xfemal, d1=dfemal, y1=yfemal,
  x2=xmale, d2=dmale, y2=ymale, theta=c(0.25,0), fun1=fun2, fun2=fun2)
##### you get "-2LLR" = 1.554386, etc #####
}

```

emplikHs.test2

Two sample empirical likelihood ratio test for hazards with right censored, left truncated data. Many constraints.

Description

Use empirical likelihood ratio and Wilks theorem to test the null hypothesis that

$$\int f_1(t)dH_1(t) - \int f_2(t)dH_2(t) = \theta$$

where $H_*(t)$ is the (unknown) cumulative hazard functions; $f_*(t)$ can be any predictable functions of t . θ is a vector of parameters (dim=q). The given value of θ in these computation are the value to be tested. The data can be right censored and left truncated.

When the given constants θ is too far away from the NPMLE, there will be no hazard function satisfy this constraint and the -2 Log empirical likelihood ratio will be infinite. In this case the computation will stop.

Usage

```
emplikHs.test2(x1, d1, y1= -Inf, x2, d2, y2 = -Inf,
  theta, fun1, fun2, maxit=25,tola = 1e-7,itertrace =FALSE)
```

Arguments

x1 a vector of length n1, the observed survival times, sample 1.
d1 a vector, the censoring indicators, 1-uncensor; 0-censor.
y1 optional vector, the left truncation times.
x2 a vector of length n2, the observed survival times, sample 2.

d2	a vector, the censoring indicators, 1-uncensor; 0-censor.
y2	optional vector, the left truncation times.
fun1	a predictable function used to calculate the weighted discrete hazard to form the null hypothesis H_0 . fun1(x) must be able to take a vector input (length n1) x, and output a matrix of n1 x q. When q=1, the output can also be a vector.
fun2	Ditto. but for length n2
tola	an optional positive real number, the tolerance of iteration error in solve the non-linear equation needed in constrained maximization.
theta	a given vector of length q. for Ho constraint.
maxit	integer, maximum number of Newton-Raphson type iterations.
itertrace	Logical, if the results of each iteration needs to be printed.

Details

The log likelihood been maximized is the Poisson likelihood:

$$\sum D_{1i} \log w_i - \sum R_{1i} w_i + \sum D_{2j} \log v_j - \sum R_{2j} v_j$$

where $w_i = \Delta H_1(t_i)$ is the jump of the cumulative hazard function at t_i (for first sample), D_{1i} is the number of failures observed at t_i , R_{1i} is the number of subjects at risk at time t_i . Dido for sample two.

For (proper) discrete distributions, the jump size of the cumulative hazard at the last jump is always 1. So, in the likelihood ratio, it cancels. But the last jump of size 1 still matter when computing the constraint.

The constants theta must be inside the so called feasible region for the computation to continue. This is similar to the requirement that in testing the value of the mean, the value must be inside the convex hull of the observations. It is always true that the NPMLE values are feasible. So when the computation stops, try move the theta closer to the NPMLE, which we print out first thing in this function, even when other later computations do not go. When the computation stops, the -2LLR should have value infinite.

This function uses the llog etc. function, so sometimes it may produce different result from the one sample result. which use the regular log function. The advantage is that we avoid the possible log(0) situation.

You can also use this function for one sample problems. You need to artificially supply data for sample two of minimal size (like size 2q+2), and specify a fun2() that ALWAYS return 0's (zero vector, with length=n2 vector length, or zero matrix, with dim n2 x q as the input). Then, look for -2LLR(sample1) in the output.

Value

A list with the following components:

"-2LLR"	The -2Log empirical Likelihood ratio.
lambda	the final value of the Lagrange multiplier.

```
"-2LLR(sample1)"
```

The -2Log empirical likelihood ratio for sample one only. Useful in one sample problems.

```
"Llog(sample1)"
```

The numerator only of the above "-2LLR(sample1)", without -2.

Author(s)

Mai Zhou

References

Zhou and Fang (2001). "Empirical likelihood ratio for 2 sample problems for censored data". *Tech Report, Univ. of Kentucky, Dept of Statistics*

See Also

emplikH2.test

Examples

```
if(require("boot", quietly = TRUE)) {
  ###library(boot)
  data(channing)
  ymale <- channing[1:97,2]
  dmale <- channing[1:97,5]
  xmale <- channing[1:97,3]
  yfemale <- channing[98:462,2]
  dfemale <- channing[98:462,5]
  xfemale <- channing[98:462,3]
  fun1 <- function(x) { as.numeric(x <= 960) }
  #####
  fun2 <- function(x){ cbind(as.numeric(x <= 960), as.numeric(x <= 860))}
  ##### fun2 has matrix output #####
  emplikHs.test2(x1=xfemale, d1=dfemale, y1=yfemale,
    x2=xmale, d2=dmale, y2=ymale, theta=c(0,0), fun1=fun2, fun2=fun2)
}
#####
##### Second example:
if(require("KMsurv", quietly = TRUE)) {
  ###library(KMsurv)
  data(kidney)
  ### these functions counts the risk set size, so delta=1 always ###
  temp1 <- Wdataclean3(z=kidney$time[kidney[,3]==1], d=rep(1,43) )
  temp2 <- DnR(x=temp1$value, d=temp1$dd, w=temp1$weight)
  TIME <- temp2$times
  RISK <- temp2$n.risk
  fR1 <- approxfun(x=TIME, y=RISK, method="constant", yright=0, rule=2, f=1)
  temp1 <- Wdataclean3(z=kidney$time[kidney[,3]==2], d=rep(1,76) )
  temp2 <- DnR(x=temp1$value, d=temp1$dd, w=temp1$weight)
  TIME <- temp2$times
  RISK <- temp2$n.risk
```

```

fR2 <- approxfun(x=TIME, y=RISK, method="constant", yright=0, rule=2, f=1)

### the weight function for two sample Gehan-Wilcoxon type test ###
fun <- function(t){ fR1(t)*fR2(t)/((76*43)*sqrt(119/(76*43))) }
### Here comes the test: ###
emplikHs.test2(x1=kidney[kidney[,3]==1,1],d1=kidney[kidney[,3]==1,2],
  x2=kidney[kidney[,3]==2,1],d2=kidney[kidney[,3]==2,2],
  theta=0, fun1= fun, fun2=fun)
### The results should include this ###
#"$"-2LLR"
#[1] 0.002473070
#
#$lambda
#[1] -0.1713749
#####
##### the weight function for log-rank test #####
funlogrank <- function(t){sqrt(119/(76*43))*fR1(t)*fR2(t)/(fR1(t)+fR2(t))}
##### Now the log-rank test ###
emplikHs.test2(x1=kidney[kidney[,3]==1,1],d1=kidney[kidney[,3]==1,2],
  x2=kidney[kidney[,3]==2,1],d2=kidney[kidney[,3]==2,2],
  theta=0, fun1=funlogrank, fun2=funlogrank)
##### The result of log rank test should include this ###
#
#"$"-2LLR"
#[1] 2.655808
#
#$lambda
#[1] 3.568833
#####
##### the weight function for both type test #####
funBOTH <- function(t) {
  cbind(sqrt(119/(76*43))*fR1(t)*fR2(t)/(fR1(t)+fR2(t)),
    fR1(t)*fR2(t)/((76*43)*sqrt(119/(76*43)))) }
##### The test that combine both tests ###
emplikHs.test2(x1=kidney[kidney$type==1,1],d1=kidney[kidney$type==1,2],
  x2=kidney[kidney$type==2,1],d2=kidney[kidney$type==2,2],
  theta=c(0,0), fun1=funBOTH, fun2=funBOTH)
##### the result should include this ###
#
#"$"-2LLR"
#[1] 13.25476
#
#$lambda
#[1] 14.80228 -21.86733
#####
}

```


Description

This function is similar to `findUL2` but here the seeking of lower and upper bound are separate (the other half is `findUnew`).

The reason for this is: sometime we need to supply the fun with different nuisance parameter(s) values when seeking Lower or Upper bound. For example fun returns the $-2LLR$ for a given parameter of interest, but there are additional nuisance parameter need to be profiled out, and we need to give a range of the nuisance parameter to be max/min over. This range can be very different for parameter near Upper bound vs near Lower bound. In the `findUL2`, you have to supply a range really wide that (hopefully) works for both Upper and Lower bound. Here, with separate `findLnew` and `findUnew` we can tailor the range for one end of the confidence interval.

Those nuisance parameter(s) are supplied via the ... input of this function.

Another improvement is that we used the "extendInt" option of the `uniroot`. So now we can and did used a smaller default step input, compare to `findUL2`.

This program uses `uniroot()` to find (only) the lower (Wilks) confidence limit based on the -2 log likelihood ratio, which the required input fun is supposed to supply.

Basically, starting from MLE, we search on lower direction, by step away from MLE, until we find values that have $-2LLR = level$. (the value of $-2LLR$ at MLE is supposed to be zero.)

At current implementation, only handles one dimensional parameter, i.e. only confidence intervals, not confidence regions.

Usage

```
findLnew(step=0.003, initStep=0, fun, MLE, level=3.84146, tol=.Machine$double.eps^0.5, ...)
```

Arguments

<code>step</code>	a positive number. The starting step size of the search. Reasonable value should be about 1/5 of the SD of MLE.
<code>initStep</code>	a nonnegative number. The first step size of the search. Sometimes, you may want to put a larger <code>initStep</code> to speed the search.
<code>fun</code>	a function that returns a list. One of the item in the list should be " $-2LLR$ ", which is the -2 log (empirical) likelihood ratio. The first input of fun must be the parameter for which we are seeking the confidence interval. (The MLE or NPMLE of this parameter should be supplied as in the input MLE). The rest of the input to fun are typically the data. If the first input of fun is set to MLE, then the returned $-2LLR$ should be 0.
<code>MLE</code>	The MLE of the parameter. No need to be exact, as long as it is inside the confidence interval.
<code>level</code>	an optional positive number, controls the confidence level. Default to 3.84146 = <code>chisq(0.95, df=1)</code> . Change to 2.70= <code>chisq(0.90, df=1)</code> to get a 90% confidence interval.
<code>tol</code>	Error bound of the final result.
<code>...</code>	additional arguments, if any, to pass to fun.

Details

Basically we repeatedly testing the value of the parameter, until we find those which the -2 log likelihood value is equal to 3.84146 (or other level, if set differently).

Value

A list with the following components:

Low	the lower limit of the confidence interval.
FstepL	the final step size when search lower limit. An indication of the precision.
Lvalue	The -2LLR value of the final Low value. Should be approximately equal to level. If larger than level, than the confidence interval limit Low is wrong.

Author(s)

Mai Zhou

References

Zhou, M. (2016). Empirical Likelihood Method in Survival Analysis. CRC Press.

Examples

```
## example with tied observations. Kaplan-Meier mean=4.0659.
## For more examples see vignettes.
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
myfun6 <- function(theta, x, d) {
  e1.cen.EM2(x, d, fun=function(t){t}, mu=theta)
}
findLnew(step=0.1, fun=myfun6, MLE=4.0659, x=x, d=d)
```

findUL

Find the Wilks Confidence Interval from the Given (empirical) Likelihood Ratio Function

Description

This program uses `uniroot()` to find the upper and lower (Wilks) confidence limits based on the -2 log likelihood ratio, which the required input `fun` is supposed to supply.

Basically, starting from MLE, we search on both directions, by `step` away from MLE, until we find values that have $-2LLR = \text{level}$. (the value of $-2LLR$ at MLE is supposed to be zero.)

At curruent implimentation, only handles one dimesional parameter, i.e. only confidence intervals, not confidence regions.

For examples of using this function to find confidence interval, see the pdf vignettes file.

Usage

```
findUL (step = 0.01, initStep = 0, fun, MLE, level = 3.84146, ...)
```

Arguments

step	a positive number. The starting step size of the search. Reasonable value should be about 1/5 of the SD of MLE.
initStep	a nonnegative number. The first step size of the search. Sometimes, you may want to put a larger innitStep to speed the search.
fun	a function that returns a list. One of the item in the list should be "-2LLR", which is the -2 log (empirical) likelihood ratio. The first input of fun must be the parameter for which we are seeking the confidence interval. (The MLE or NPMLE of this parameter should be supplied as in the input MLE). The rest of the input to fun are typically the data. If the first input of fun is set to MLE, then the returned -2LLR should be 0.
MLE	The MLE of the parameter. No need to be exact, as long as it is inside the confidence interval.
level	an optional positive number, controls the confidence level. Default to 3.84 = $\text{chisq}(0.95, \text{df}=1)$. Change to 2.70= $\text{chisq}(0.90, \text{df}=1)$ to get a 90% confidence interval.
...	additional arguments, if any, to pass to fun.

Details

Basically we repeatedly testing the value of the parameter, until we find those which the -2 log likelihood value is equal to 3.84 (or other level, if set differently).

If there is no value exactly equal to 3.84, it is better to use findULold(), where we stop at the value which result a -2 log likelihood just below 3.84. (as in the discrete case, like quantiles.)

Value

A list with the following components:

Low	the lower limit of the confidence interval.
Up	the upper limit of the confidence interval.
FstepL	the final step size when search lower limit. An indication of the precision.
FstepU	Ditto. An indication of the precision of the upper limit.
Lvalue	The -2LLR value of the final Low value. Should be approximately equal to level. If larger than level, than the confidence interval limit Low is wrong.
Uvalue	Ditto. Should be approximately equa to level.

Author(s)

Mai Zhou

References

Zhou, M. (2016). Empirical Likelihood Method in Survival Analysis. CRC Press.

Examples

```
## example with tied observations. Kaplan-Meier mean=4.0659.
## For more examples see vignettes.
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
myfun6 <- function(theta, x, d) {
  el.cen.EM2(x, d, fun=function(t){t}, mu=theta)
}
findUL(step=0.2, fun=myfun6, MLE=4.0659, x=x, d=d)
```

findUL2	<i>Find the Wilks Confidence Interval from the Given (empirical) Likelihood Ratio Function</i>
---------	--

Description

This program uses simple search and uniroot() to find the upper and lower (Wilks) confidence limits based on the $-2 \log$ likelihood ratio, which the required input fun is supposed to supply.

This function is faster than findUL().

Basically, starting from MLE, we search on both directions, by step away from MLE, until we find values that have $-2LLR = \text{level}$. (the value of $-2LLR$ at MLE is supposed to be zero.)

At current implementation, only handles one dimensional parameter, i.e. only confidence intervals, not confidence regions.

For examples of using this function to find confidence interval, see the pdf vignettes file.

Usage

```
findUL2(step=0.01, initStep=0, fun, MLE, level=3.84146, tol=.Machine$double.eps^0.5, ...)
```

Arguments

step	a positive number. The starting step size of the search. Reasonable value should be about 1/5 of the SD of MLE.
initStep	a nonnegative number. The first step size of the search. Sometimes, you may want to put a larger innitStep to speed the search.
fun	a function that returns a list. One of the item in the list should be "-2LLR", which is the $-2 \log$ (empirical) likelihood ratio. The first input of fun must be the parameter for which we are seeking the confidence interval. (The MLE or NPMLE of this parameter should be supplied as in the input MLE). The rest of the input to fun are typically the data. If the first input of fun is set to MLE, then the returned $-2LLR$ should be 0.

MLE	The MLE of the parameter. No need to be exact, as long as it is inside the confidence interval.
level	an optional positive number, controls the confidence level. Default to 3.84 = $\text{chisq}(0.95, \text{df}=1)$. Change to 2.70= $\text{chisq}(0.90, \text{df}=1)$ to get a 90% confidence interval.
tol	tolerance to pass to <code>uniroot()</code> . Default to <code>.Machine\$double.eps^0.5</code>
...	additional arguments, if any, to pass to <code>fun</code> .

Details

Basically we repeatedly testing the value of the parameter, until we find those which the -2 log likelihood value is equal to 3.84 (or other level, if set differently).

If there is no value exactly equal to 3.84, we stop at the value which result a -2 log likelihood just below 3.84. (as in the discrete case, like quantiles.)

Value

A list with the following components:

Low	the lower limit of the confidence interval.
Up	the upper limit of the confidence interval.
FstepL	the final step size when search lower limit. An indication of the precision.
FstepU	Ditto. An indication of the precision of the upper limit.
Lvalue	The -2LLR value of the final Low value. Should be approximately equal to level. If larger than level, than the confidence interval limit Low is wrong.
Uvalue	Ditto. Should be approximately equa to level.

Author(s)

Mai Zhou

References

Zhou, M. (2016). Empirical Likelihood Method in Survival Analysis. CRC Press.

Zhou, M. (2002). Computing censored empirical likelihood ratio by EM algorithm. *JCGS*

Examples

```
## example with tied observations. Kaplan-Meier mean=4.0659.
## For more examples see vignettes.
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
myfun6 <- function(theta, x, d) {
  el.cen.EM2(x, d, fun=function(t){t}, mu=theta)
}
findUL2(step=0.2, fun=myfun6, MLE=4.0659, x=x, d=d)
```

 findULold

Find the Wilks Confidence Interval from the Given (empirical) Likelihood Ratio Function

Description

This program uses simple search to find the upper and lower (Wilks) confidence limits based on the -2 log likelihood ratio, which the required input fun is supposed to supply.

Basically, starting from MLE, we search on both directions, by step away from MLE, until we find values that have $-2LLR = \text{level}$. (the value of $-2LLR$ at MLE is supposed to be zero.)

At current implementation, only handles one dimensional parameter, i.e. only confidence intervals, not confidence regions.

For examples of using this function to find confidence interval, see the pdf vignettes file.

Usage

```
findULold (step = 0.01, initStep = 0, fun, MLE, level = 3.84146, ...)
```

Arguments

step	a positive number. The starting step size of the search. Reasonable value should be about 1/5 of the SD of MLE.
initStep	a nonnegative number. The first step size of the search. Sometimes, you may want to put a larger innitStep to speed the search.
fun	a function that returns a list. One of the item in the list should be "-2LLR", which is the -2 log (empirical) likelihood ratio. The first input of fun must be the parameter for which we are seeking the confidence interval. (The MLE or NPMLE of this parameter should be supplied as in the input MLE). The rest of the input to fun are typically the data. If the first input of fun is set to MLE, then the returned $-2LLR$ should be 0.
MLE	The MLE of the parameter. No need to be exact, as long as it is inside the confidence interval.
level	an optional positive number, controls the confidence level. Default to 3.84146 = $\text{chisq}(0.95, \text{df}=1)$. Change to 2.70= $\text{chisq}(0.90, \text{df}=1)$ to get a 90% confidence interval.
...	additional arguments, if any, to pass to fun.

Details

Basically we repeatedly testing the value of the parameter, until we find those which the -2 log likelihood value is equal to 3.84146 (or other level, if set differently).

If there is no value exactly equal to 3.84146, we stop at the value which result a -2 log likelihood just below 3.84146. (as in the discrete case, like quantiles.)

Value

A list with the following components:

Low	the lower limit of the confidence interval.
Up	the upper limit of the confidence interval.
FstepL	the final step size when search lower limit. An indication of the precision.
FstepU	Ditto. An indication of the precision of the upper limit.
Lvalue	The -2LLR value of the final Low value. Should be approximately equal to level. If larger than level, than the confidence interval limit Low is wrong.
Uvalue	Ditto. Should be approximately equa to level.

Author(s)

Mai Zhou

References

Zhou, M. (2016). Empirical Likelihood Method in Survival Analysis. CRC Press.

Examples

```
## example with tied observations. Kaplan-Meier mean=4.0659.
## For more examples see vignettes.
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
myfun6 <- function(theta, x, d) {
  el.cen.EM2(x, d, fun=function(t){t}, mu=theta)
}
## findULold(step=0.2, fun=myfun6, MLE=4.0659, level = qchisq(0.9, df=1) , x=x, d=d)
```

findUnew

Find the Wilks Confidence Interval Upper Bound from the Given (empirical) Likelihood Ratio Function

Description

This function is similar to findUL2 but here the seeking of lower and upper bound are separate (the other half is findLnew).

See the help file of findLnew. Since sometimes the likelihood ratio is a Profile likelihood ratio and we need to supply the fun with different nuisance parameter(s) value(s) when seeking Lower or Upper bound. Those nuisance parameter(s) are supplied via the ... input.

Another improvement is that we used the "extendInt" option of the uniroot. So now we can and did used a smaller default step input, 0.003, compare to findUL2.

This program uses uniroot() to find (only) the upper (Wilks) confidence limit based on the -2 log likelihood ratio, which the required input fun is supposed to supply.

Basically, starting from MLE, we search on upper/higher direction, by step away from MLE, until we find values that have $-2LLR = \text{level}$. (the value of $-2LLR$ at MLE is supposed to be zero.)

At current implementation, only handles one dimensional parameter, i.e. only confidence intervals, not confidence regions.

Usage

```
findUnew(step=0.003, initStep=0, fun, MLE, level=3.84146, tol=.Machine$double.eps^0.5, ...)
```

Arguments

step	a positive number. The starting step size of the search. Reasonable value should be about 1/5 of the SD of MLE.
initStep	a nonnegative number. The first step size of the search. Sometimes, you may want to put a larger inittStep to speed the search.
fun	a function that returns a list. One of the item in the list should be "-2LLR", which is the $-2 \log$ (empirical) likelihood ratio. The first input of fun must be the parameter for which we are seeking the confidence interval. (The MLE or NPMLE of this parameter should be supplied as in the input MLE). The rest of the input to fun are typically the data. If the first input of fun is set to MLE, then the returned $-2LLR$ should be 0.
MLE	The MLE of the parameter. No need to be exact, as long as it is inside the confidence interval.
level	an optional positive number, controls the confidence level. Default to 3.84146 = $\text{chisq}(0.95, \text{df}=1)$. Change to 2.70= $\text{chisq}(0.90, \text{df}=1)$ to get a 90% confidence interval.
tol	Error bound of the final result.
...	additional arguments, if any, to pass to fun.

Details

Basically we repeatedly testing the value of the parameter, until we find those which the $-2 \log$ likelihood value is equal to 3.84146 (or other level, if set differently).

Value

A list with the following components:

Up	the lower limit of the confidence interval.
FstepU	the final step size when search lower limit. An indication of the precision/error size.
Uvalue	The $-2LLR$ value of the final Up value. Should be approximately equal to level. If larger than level, than the confidence interval limit Up is wrong.

Author(s)

Mai Zhou

References

Zhou, M. (2016). Empirical Likelihood Method in Survival Analysis. CRC Press.

Examples

```
## example with tied observations. Kaplan-Meier mean=4.0659.
## For more examples see vignettes.
x <- c(1, 1.5, 2, 3, 4, 5, 6, 5, 4, 1, 2, 4.5)
d <- c(1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 1)
myfun6 <- function(theta, x, d) {
  el.cen.EM2(x, d, fun=function(t){t}, mu=theta)
}
findUnew(step=0.1, fun=myfun6, MLE=4.0659, x=x, d=d)
```

myeloma

Multiple Myeloma Data

Description

Krall, Uthoff, and Harley (1975) analyzed data from a study on multiple myeloma in which researchers treated 65 patients with alkylating agents.

Of those patients, 48 died during the study and 17 survived. In the data set MYELOMA, the variable TIME represents the survival time in months from diagnosis. The variable VSTATUS consists of two values, 0 and 1, indicating whether the patient was alive or dead, respectively, at the of end the study. If the value of VSTATUS is 0, the corresponding value of TIME is censored.

The variables thought to be related to survival are LOGBUN (log BUN at diagnosis), HGB (hemoglobin at diagnosis), PLATELET (platelets at diagnosis: 0=abnormal, 1=normal), AGE (age at diagnosis in years), LOGWBC (log WBC at diagnosis), FRAC (fractures at diagnosis: 0=none, 1=present), LOGPBM (log percentage of plasma cells in bone marrow), PROTEIN (proteinuria at diagnosis), and SCALC (serum calcium at diagnosis).

Data are from http://ftp.sas.com/techsup/download/sample/samp_lib/statsampExamples_of_Coxs_Model.html

Usage

```
data(myeloma)
```

Format

A data frame containing 65 observations on 11 variables:

```
[,1] "time"
[,2] "vstatus"
[,3] "logBUN"
[,4] "HGB"
[,5] "platelet"
```

[,6] "age"
 [,7] "logWBC"
 [,8] "FRAC"
 [,9] "logPBM"
 [,10] "protein"
 [,11] "SCALC"

References

Krall, J.M., Uthoff, V.A., and Harley, J. B. (1975). A Step-up Procedure for Selecting Variables Associated with Survival. *Biometrics*, 31, 49-57.

RankRegTest	<i>Test the AFT model Rank Regression estimator by Empirical Likelihood</i>
-------------	---

Description

Use the empirical likelihood ratio and Wilks theorem to test if the regression coefficient is equal to beta, based on the rank estimator for the AFT model.

The log empirical likelihood been maximized is

$$\sum_{d=1} \log \Delta F(e_i) + \sum_{d=0} \log[1 - F(e_i)];$$

where e_i are the residuals.

Usage

```
RankRegTest(y, d, x, beta, type="Gehan")
```

Arguments

y	a vector of length N, containing the censored responses.
d	a vector (length N) of either 1's or 0's. d=1 means y is uncensored; d=0 means y is right censored.
x	a matrix of size N by q.
beta	a vector of length q. the value of the regression coefficient to be tested in the model $y_i = \beta x_i + \epsilon_i$.
type	default to Gehan type. The other option is Logrank type.

Details

The estimator of beta can be obtained by function `rankaft()` in the package `rankreg`. But here you may test other values of beta. If you test the beta value that is obtained from the `rankaft()`, then the -2LLR should be 0 and the p-value should be 1.

The above likelihood should be understood as the likelihood of the error term, so in the regression model the error e_i should be iid.

The estimation equation used when maximize the empirical likelihood is

$$0 = \sum_i \phi(e_i) d_i \Delta F(e_i) (x_i - \bar{x}_i) / (nw_i)$$

which was described in detail in the references below.

See also the function `RankRegTestH`, which is based on the hazard likelihood.

Value

A list with the following components:

"-2LLR"	the -2 loglikelihood ratio; should have approximate chisq distribution under H_0 .
loge12	the log empirical likelihood, under estimating equation.
loge1	the log empirical likelihood of the Kaplan-Meier of e's.
prob	the probabilities that max the empirical likelihood under rank estimating equation constraint.

Author(s)

Mai Zhou.

References

Kalbfleisch, J. and Prentice, R. (2002) *The Statistical Analysis of Failure Time Data*. 2nd Ed. Wiley, New York. (Chapter 7)

Jin, Z., Lin, D.Y., Wei, L. J. and Ying, Z. (2003). Rank-based inference for the accelerated failure time model. *Biometrika*, **90**, 341-53.

Zhou, M. (2005). Empirical likelihood analysis of the rank estimator for the censored accelerated failure time model. *Biometrika*, **92**, 492-98.

Examples

```
data(myeloma)
RankRegTest(y=myeloma[,1], d=myeloma[,2], x=myeloma[,3], beta= -15.50147)
# you should get "-2LLR" = 9.050426e-05 (practically zero)
# The beta value, -15.50147, was obtained by rankaft() from the rankreg package.
```

RankRegTestH	<i>Test the AFT model, Rank Regression estimator by (Hazard)Empirical Likelihood</i>
--------------	--

Description

Use the empirical likelihood ratio and Wilks theorem to test if the regression coefficient is equal to beta, based on the rank estimator/estimating equation of the AFT model.

The log empirical likelihood been maximized is the hazard empirical likelihood.

Usage

```
RankRegTestH(y, d, x, beta, type="Gehan")
```

Arguments

y	a vector of length N, containing the censored responses.
d	a vector (length N) of either 1's or 0's. d=1 means y is uncensored; d=0 means y is right censored.
x	a matrix of size N by q.
beta	a vector of length q. the value of the regression coefficient to be tested in the model $y_i = \beta x_i + \epsilon_i$.
type	default to Gehan type. The other option is Logrank type.

Details

The estimator of beta can be obtained by function `rankaft()` in the package `rankreg`. But here you may test other values of beta. If you test the beta value that is obtained from the `rankaft()`, then the -2LLR should be 0 and the p-value should be 1.

The above likelihood should be understood as the likelihood of the error term, so in the regression model the error e_i should be iid.

The estimating equation used when maximize the empirical likelihood is

$$0 = \sum_i R(e_i) \phi(e_i) d_i \Delta A(e_i) (x_i - \bar{x}_i)$$

where all notation was described in detail in the references below.

Value

A list with the following components:

"-2LLR"	the -2 loglikelihood ratio; should have approximate chisq distribution under H_0 .
loge12	the log empirical likelihood, under estimating equation.
loge1	the log empirical likelihood of the Kaplan-Meier of e's.

Author(s)

Mai Zhou

References

- Zhou, M. (2016) Empirical Likelihood Methods in Survival Analysis. CRC Press.
- Kalbfleisch, J. and Prentice, R. (2002) The Statistical Analysis of Failure Time Data. 2nd Ed. Wiley, New York. (Chapter 7)
- Jin, Z., Lin, D.Y., Wei, L. J. and Ying, Z. (2003). Rank-based inference for the accelerated failure time model. *Biometrika*, **90**, 341-53.
- Zhou, M. (2005). Empirical likelihood analysis of the rank estimator for the censored accelerated failure time model. *Biometrika*, **92**, 492–498.

Examples

```
data(myeloma)
RankRegTestH(y=myeloma[,1], d=myeloma[,2], x=myeloma[,3], beta= -15.50147)
# you should get "-2LLR" = 9.050426e-05 (practically zero)
# The beta value, -15.50147, was obtained by rankaft() from
# the rankreg package.
```

ROCnp

*Test the ROC curve by Empirical Likelihood***Description**

Use empirical likelihood ratio to test the hypothesis H_0 : $(1-b_0)$ th quantile of sample 1 = $(1-t_0)$ th quantile of sample 2. This is the same as testing H_0 : $R(t_0) = b_0$, where $R(\cdot)$ is the ROC curve.

The log empirical likelihood been maximized is

$$\sum_{d_1=1} \log \Delta F_1(t_{1i}) + \sum_{d_1=0} \log[1 - F_1(t_{1i})] + \sum_{d_2=1} \log \Delta F_2(t_{2j}) + \sum_{d_2=0} \log[1 - F_2(t_{2j})].$$

This empirical likelihood ratio has a chi square limit under H_0 .

Usage

```
ROCnp(t1, d1, t2, d2, b0, t0)
```

Arguments

- | | |
|----|---|
| t1 | a vector of length n. Observed times, may be right censored. |
| d1 | a vector of length n, censoring status. d=1 means t is uncensored; d=0 means t is right censored. |
| t2 | a vector of length m. Observed times, may be right censored. |
| d2 | a vector of length m, censoring status. |
| b0 | a scalar between 0 and 1. |
| t0 | a scalar, betwenn 0 and 1. |

Details

Basically, we first test $(1-b_0)$ th quantile of sample 1 = c and also test $(1-t_0)$ th quantile of sample 2 = c . This way we obtain two log likelihood ratios.

Then we minimize the sum of the two log likelihood ratio over c .

See the tech report below for details on a similar setting.

Value

A list with the following components:

"-2LLR" the -2 loglikelihood ratio; have approximate chisq distribution under H_0 .
 cstar the estimated common quantile.

Author(s)

Mai Zhou.

References

Zhou, M. and Liang, H (2008). Empirical Likelihood for Hybrid Two Sample Problem with Censored Data. *Univ. Kentucky Tech. Report*.

Su, H., Zhou, M. and Liang, H. (2011). Semi-parametric hybrid empirical likelihood inference for two-sample comparison with censored data. *Lifetime Data Analysis*, **17**, 533-551.

Examples

```
#### An example of testing the equality of two medians. No censoring.
ROCnp(t1=rexp(100), d1=rep(1,100), t2=rexp(120), d2=rep(1,120), b0=0.5, t0=0.5)
#####
#### Next, an example of finding 90 percent confidence interval of R(0.5)
#### Note: We are finding confidence interval for R(0.5). So we are testing
#### R(0.5)= 0.35, 0.36, 0.37, 0.38, etc. try to find values so that
#### testing R(0.5) = L , U has p-value of 0.10, then [L, U] is the 90 percent
#### confidence interval for R(0.5).
#set.seed(123)
#t1 <- rexp(200)
#t2 <- rexp(200)
#ROCnp( t1=t1, d1=rep(1, 200), t2=t2, d2=rep(1, 200), b0=0.5, t0=0.5)$"-2LLR"
#### since the -2LLR value is less than 2.705543 = qchisq(0.9, df=1), so the
#### confidence interval contains 0.5.
#gridpoints <- 35:65/100
#ELvalues <- gridpoints
#for( i in 1:31 ) ELvalues[i] <- ROCnp(t1=t1, d1=rep(1, 200),
#   t2=t2, d2=rep(1, 200), b0=gridpoints[i], t0=0.5)$"-2LLR"
#myfun1 <- approxfun(x=gridpoints, y=ELvalues)
#uniroot( f= function(x){myfun1(x)-2.705543}, interval= c(0.35, 0.5) )
#uniroot( f= function(x){myfun1(x)-2.705543}, interval= c(0.5, 0.65) )
#### So, taking the two roots, we see the 90 percent confidence interval for R(0.5)
#### in this case is [0.4478081, 0.5889425].
```

Description

Use empirical likelihood ratio to test the hypothesis H_0 : $(1-b_0)$ th quantile of sample 1 = $(1-t_0)$ th quantile of sample 2. This is the same as testing H_0 : $R(t_0) = b_0$, where $R(\cdot)$ is the ROC curve.

The log empirical likelihood been maximized is

$$\sum_{d1=1} \log \Delta F_1(t1_i) + \sum_{d1=0} \log[1 - F_1(t1_i)] + \sum_{d2=1} \log \Delta F_2(t2_j) + \sum_{d2=0} \log[1 - F_2(t2_j)].$$

This empirical likelihood ratio has a chi square limit under H_0 .

Usage

ROCnp2(t1, d1, t2, d2, b0, t0)

Arguments

t1	a vector of length n. Observed times, sample 1. may be right censored.
d1	a vector of length n, censoring status. d=1 means t is uncensored; d=0 means t is right censored.
t2	a vector of length m. Observed times, sample 2. may be right censored.
d2	a vector of length m, censoring status.
b0	a scalar, between 0 and 1.
t0	a scalar, between 0 and 1.

Details

First, we test $(1-b_0)$ th quantile of sample 1 = c and also test $(1-t_0)$ th quantile of sample 2 = c . This way we obtain two log likelihood ratios.

Then we minimize the sum of the two log likelihood ratios over c .

This version use an exhaust search for the minimum (over c). Since the objective (log lik) are piecewise constants, the optimum() function in R do not work well. See the tech report below for details on a similar setting.

Value

A list with the following components:

"-2LLR"	the -2 loglikelihood ratio; have approximate chisq distribution under H_0 .
cstar	the estimated common quantile.

Author(s)

Mai Zhou

References

Su, Haiyan; Zhou, Mai and Liang, Hua (2011). Semi-parametric Hybrid Empirical Likelihood Inference for Two Sample Comparison with Censored Data. *Lifetime Data Analysis*, **17**, 533-551.

Examples

```
##### An example of testing the equality of two medians.
##### No censoring.
# ROCnp2(t1=rexp(100), d1=rep(1,100), t2=rexp(120),
#           d2=rep(1,120), b0=0.5, t0=0.5)
#####
##### This example do not work on the Solaris Sparc machine.
##### But works fine on other platforms.
#####
##### Next, an example of finding 90 percent confidence
##### interval of R(0.5)
##### Note: We are finding confidence interval for R(0.5).
##### So we are testing
##### R(0.5)= 0.35, 0.36, 0.37, 0.38, etc. try to find
##### values so that testing R(0.5) = L , U has p-value
##### of 0.10, then [L, U] is the 90 percent
##### confidence interval for R(0.5).
#set.seed(123)
#t1 <- rexp(200)
#t2 <- rexp(200)
#ROCnp( t1=t1, d1=rep(1, 200), t2=t2, d2=rep(1, 200),
#       b0=0.5, t0=0.5)$"-2LLR"
##### since the -2LLR value is less than
##### 2.705543 = qchisq(0.9, df=1), so the
##### confidence interval contains 0.5.
#gridpoints <- 35:65/100
#ELvalues <- gridpoints
#for(i in 1:31) ELvalues[i] <- ROCnp2(t1=t1, d1=rep(1, 200),
#   t2=t2, d2=rep(1, 200), b0=gridpoints[i], t0=0.5)$"-2LLR"
#myfun1 <- approxfun(x=gridpoints, y=ELvalues)
#uniroot(f=function(x){myfun1(x)-2.705543},
#        interval= c(0.35, 0.5) )
#uniroot(f= function(x){myfun1(x)-2.705543},
#        interval= c(0.5, 0.65) )
##### So, taking the two roots, we see the 90 percent
##### confidence interval for R(0.5) in this
##### case is [0.4457862, 0.5907723].
#####
```

`smallcell`*Smallcell Lung Cancer Data*

Description

There are 121 observations on 4 variables. Arm is the indication of two treatments. (either C + E or E + C). Entry is the age of the patient at entry. Survival is the survival time and indicator is the censoring indicator (right censoring). For more details please see the reference below.

Data are from Ying, Z., Jung, SH, and Wei, LJ (1995). Median regression analysis with censored data. *Journal of the American Statistical Association*, 90, 178-184.

Usage

```
data(smallcell)
```

Format

A data frame containing 121 observations on 4 variables:

```
[,1] "arm"  
[,2] "entry"  
[,3] "survival"  
[,4] "indicator"
```

References

Ying, Z., Jung, SH, and Wei, LJ (1995). Median regression analysis with censored data. *Journal of the American Statistical Association*, 90, 178-184.

Maksymiuk, A. W., Jett, J. R., Earle, J. D., Su, J. Q., Diegert, F. A., Mailliard, J. A., Kardinal, C. G., Krook, J. E., Veeder, M. H., Wiesenfeld, M., Tschetter, L. K., and Levitt, R. (1994). Sequencing and Schedule Effects of Cisplatin Plus Etoposide in Small Cell Lung Cancer Results of a North Central Cancer Treatment Group Randomized Clinical Trial. *Journal of Clinical Oncology*, 12, 70-76.

`WRegEst`*Compute the casewise weighted regression estimator for AFT model*

Description

For the AFT model, this function computes the case weighted estimator of beta. Either the least squares estimator or the regression quantile estimator.

Usage

```
WRegEst(x, y, delta, LS=TRUE, tau=0.5)
```



```

WRegEst(x=cbind(1,smallcell[,1],smallcell[,2]),
        y=log10(smallcell[,3]), delta=smallcell[,4], LS=FALSE)
#####
#### you should get
####      [1]      2.603342985  -0.263000044  0.003836832
#####

```

WRegTest

Test the case weighted regression estimator by Empirical Likelihood

Description

Use the empirical likelihood ratio and Wilks theorem to test if the regression coefficient is equal to β_0 , by the case weighted estimation method.

The log empirical likelihood been maximized is

$$\sum_{d=1} \log \Delta F(y_i) + \sum_{d=0} \log[1 - F(y_i)].$$

Usage

```
WRegTest(x, y, delta, beta0, psifun=function(t){t})
```

Arguments

x	a matrix of size N by q. Random design matrix.
y	a vector of length N, containing the censored responses.
delta	a vector (length N) of either 1's or 0's. delta=1 means y is uncensored; delta=0 means y is right censored.
beta0	a vector of length q. The value of the regression coefficient to be tested in the linear model .
psifun	the estimating function. The definition of it determines the type of estimator under testing.

Details

The above likelihood should be understood as the likelihood of the censored responses y and delta.

This version can handle the model where beta is a vector (of length q).

The estimation equations used when maximize the empirical likelihood is

$$0 = \sum \delta_i \Delta F(Y_i) X_i \psi(Y_i - X_i \beta_0)$$

which was described in detail in the reference below.

For median regression (Least Absolute Deviation) estimator, you should define the psifun as +1, -1 or 0 when t is > 0 , < 0 or $= 0$.

For ordinary least squares estimator, psifun should be the identity function `psifun <- function(t)t`.

Value

A list with the following components:

"-2LLR" the -2 log likelihood ratio; have approximate chisq distribution under H_0 .
P-val the p-value using the chi-square approximation.

Author(s)

Mai Zhou.

References

Zhou, M.; Kim, M. and Bathke, A. (2012). Empirical likelihood analysis of the case weighted estimator in heteroscastic AFT model. *Statistica Sinica*, **22**, 295-316.

Examples

```
xx <- c(28, -44, 29, 30, 26, 27, 22, 23, 33, 16, 24, 29, 24, 40, 21, 31, 34, -2, 25, 19)
```

Index

* datasets

- myeloma, [57](#)
- smallcell, [65](#)

* htest

- BJjoint, [3](#)
- bjtest, [4](#)
- bjtest1d, [6](#)
- bjtestII, [7](#)
- el.cen.EM, [8](#)
- el.cen.EM2, [11](#)
- el.cen.kmc1d, [15](#)
- el.cen.test, [18](#)
- el.ltrc.EM, [20](#)
- el.test, [22](#)
- el.test.wt, [23](#)
- el.test.wt2, [25](#)
- el.trun.test, [27](#)
- emplikH.disc, [28](#)
- emplikH.disc2, [30](#)
- emplikH1.test, [32](#)
- emplikH2.test, [37](#)
- emplikHs.disc2, [43](#)
- emplikHs.test2, [45](#)
- findLnew, [48](#)
- findUL, [50](#)
- findUL2, [52](#)
- findULold, [54](#)
- findUnew, [55](#)
- RankRegTest, [58](#)
- RankRegTestH, [60](#)
- ROCnp, [61](#)
- ROCnp2, [63](#)
- WRegTest, [67](#)

* nonparametric

- BJjoint, [3](#)
- bjtest, [4](#)
- bjtest1d, [6](#)
- bjtestII, [7](#)
- el.cen.EM, [8](#)

- el.cen.EM2, [11](#)
- el.cen.kmc1d, [15](#)
- el.cen.test, [18](#)
- el.ltrc.EM, [20](#)
- el.test, [22](#)
- el.test.wt, [23](#)
- el.test.wt2, [25](#)
- el.trun.test, [27](#)
- emplikH.disc, [28](#)
- emplikH.disc2, [30](#)
- emplikH1.test, [32](#)
- emplikH1B, [34](#)
- emplikH1P, [35](#)
- emplikH2.test, [37](#)
- emplikH2B, [39](#)
- emplikH2P, [41](#)
- emplikHs.disc2, [43](#)
- emplikHs.test2, [45](#)
- findLnew, [48](#)
- findUL, [50](#)
- findUL2, [52](#)
- findULold, [54](#)
- findUnew, [55](#)
- RankRegTest, [58](#)
- RankRegTestH, [60](#)
- ROCnp, [61](#)
- ROCnp2, [63](#)
- WRegEst, [65](#)
- WRegTest, [67](#)

* package

- emplik-package, [2](#)

* survival

- el.cen.EM, [8](#)
- el.cen.EM2, [11](#)
- el.cen.kmc1d, [15](#)
- el.cen.test, [18](#)
- el.ltrc.EM, [20](#)
- el.trun.test, [27](#)
- emplikH.disc, [28](#)

- emplikH.disc2, 30
 - emplikH1.test, 32
 - emplikH1B, 34
 - emplikH1P, 35
 - emplikH2.test, 37
 - emplikH2B, 39
 - emplikH2P, 41
 - emplikHs.disc2, 43
 - emplikHs.test2, 45
- BJoint, 3
- bjtest, 4
- bjtest1d, 6
- bjtestII, 7
- e1.cen.EM, 8
- e1.cen.EM2, 11
- e1.cen.kmc1d, 15
- e1.cen.test, 18
- e1.ltrc.EM, 20
- e1.test, 22
- e1.test.wt, 23
- e1.test.wt2, 25
- e1.trun.test, 27
- emplik (emplik-package), 2
- emplik-package, 2
- emplikH.disc, 28
- emplikH.disc2, 30
- emplikH1.test, 32
- emplikH1B, 34
- emplikH1P, 35
- emplikH2.test, 37
- emplikH2B, 39
- emplikH2P, 41
- emplikHs.disc2, 43
- emplikHs.test2, 45
- findLnew, 48
- findUL, 50
- findUL2, 52
- findULold, 54
- findUnew, 55
- myeloma, 57
- RankRegTest, 58
- RankRegTestH, 60
- ROCnp, 61
- ROCnp2, 63
- smallcell, 65
- WRegEst, 65
- WRegTest, 67